

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/415, 31/505		A1	(11) International Publication Number: WO 99/59584
			(43) International Publication Date: 25 November 1999 (25.11.99)
(21) International Application Number: PCT/US99/07046		(74) Agents: MAJKA, Joseph. T. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).	
(22) International Filing Date: 17 May 1999 (17.05.99)			
(30) Priority Data: 09/081,640 20 May 1998 (20.05.98) US 09/082,977 21 May 1998 (21.05.98) US 09/106,517 29 June 1998 (29.06.98) US		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 09/081,640 (CIP) Filed on 20 May 1998 (20.05.98) US 09/082,977 (CIP) Filed on 21 May 1998 (21.05.98) US 09/106,517 (CIP) Filed on 29 June 1998 (29.06.98)		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant (for all designated States except US): SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).			
(72) Inventor; and (75) Inventor/Applicant (for US only): ESTOK, Thomas, Mark [US/US]; 1515 Charlotte Road, Plainfield, NJ 07060 (US).			
(54) Title: COMBINATION OF PHENTOLAMINE AND CYCLIC GMP PHOSPHODIESTERASE INHIBITORS FOR THE TREATMENT OF SEXUAL DYSFUNCTION			
(57) Abstract A method of treating sexual dysfunction comprising administering a therapeutically effective amount of a combination of phentolamine and cGMP PDE inhibitor such as sildenafil, as well as pharmaceutical compositions and kits useful in those methods, are disclosed.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMBINATION OF PHENTOLAMINE AND CYCLIC GMP
PHOSPHODIESTERASE INHIBITORS FOR THE TREATMENT
OF SEXUAL DYSFUNCTION

BACKGROUND

The present invention relates to pharmaceutical compositions comprising a combination of phentolamine and cyclic guanosine 3',5'-monophosphate phosphodiesterase (cGMP PDE) inhibitors and to methods of treating sexual dysfunction, especially erectile dysfunction, comprising administering an effective amount of a combination of phentolamine and cGMP PDE inhibitors.

The use of the pharmaceutical compositions and methods of this invention results in an unexpected potentiation of human sexual response.

SUMMARY OF THE INVENTION

The present invention is directed to the use of phentolamine in combination with cyclic guanosine 3',5'-monophosphate phosphodiesterase (cGMP PDE) inhibitors for the treatment of human sexual dysfunction. Preferably, the invention contemplates the use of Type V cGMP PDE inhibitor in combination with phentolamine with sildenafil being the preferred Type V cGMP PDE inhibitor.

More particularly, the present invention relates to a method of treating sexual dysfunction, especially erectile dysfunction, comprising administering to a human in need of such treatment an effective amount of a combination of phentolamine, or a pharmaceutically acceptable salt, solvate or ester thereof, and a cGMP PDE inhibitor, or a pharmaceutically acceptable salt or solvate thereof. Preferably, the invention contemplates the use of Type V cGMP PDE inhibitor in combination with phentolamine, with sildenafil being the preferred Type V cGMP PDE inhibitor.

Phentolamine mesylate and sildenafil citrate are the most preferred active ingredients for use in the methods of this invention.

In a second aspect, the invention relates to a pharmaceutical composition comprising an effective amount of phentolamine, or a pharmaceutically acceptable salt, solvate or ester thereof, and a cGMP PDE inhibitor, or a pharmaceutically acceptable salt solvate thereof. Preferably, the pharmaceutical compositions envisioned by the present invention comprise phentolamine, or a pharmaceutically acceptable salt, solvate or ester thereof, and a Type V cGMP PDE inhibitor, or a pharmaceutically acceptable salt solvate thereof, with sildenafil being the preferred Type V cGMP PDE inhibitor. Phentolamine mesylate and sildenafil citrate are the most preferred active ingredients of the pharmaceutical compositions of this invention.

In a third aspect, the invention relates to a kit comprising in one container an effective amount of phentolamine, or a pharmaceutically acceptable salt, solvate or ester thereof in a pharmaceutically acceptable carrier, and in a separate container, an effective amount of a cGMP PDE inhibitor, or a pharmaceutically acceptable salt, solvate thereof in a pharmaceutically acceptable carrier, with sildenafil being the preferred Type V cGMP PDE inhibitor. Phentolamine mesylate and sildenafil citrate are the most preferred active ingredients for use in the kits of this invention.

In a fourth aspect, the invention relates to a pharmaceutical composition for the treatment of human sexual dysfunction comprising a therapeutically effective amount of a first vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof, a therapeutically effective amount of a second vasodilating agent or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier. Preferably, the first vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof is an adrenergic blocker. More preferably, the adrenergic blocker is an alpha-adrenergic blocker. Also preferred is that the alpha adrenergic blocker is selected from the group consisting of an alpha1-adrenergic blocker, an alpha2-adrenergic blocker or both an alpha1-adrenergic blocker and an alpha2-adrenergic blocker. Preferably, the second vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof is a cGMP PDE inhibitor. Also preferred is that the first vasodilating agent or a pharmaceutically acceptable salt or solvate or

ester thereof is an adrenergic blocker and the second vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof is a cGMP PDE inhibitor. The adrenergic blocker can be selected from the group consisting of phentolamine, phentolamine mesylate, phentolamine hydrochloride, phenoxybenazmine, tolazoline, dibenamine, yohimbine, terazosin, doxazosin, prazosin and the like. The cGMP PDE inhibitor can be a cGMP PDE V inhibitor. Preferably, the cGMP PDE V inhibitor is selected from the group consisting of:

sildenafil,

(6R, 12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (Compound A), and

(3S,6R,12aR)-2,3,6,7,12,12a-hexahydro-2,3-dimethyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (Compound B) or a pharmaceutically acceptable salt or solvate thereof.

In a fifth aspect, the invention relates to a method of treating human sexual dysfunction comprising the simultaneous or sequential administration of a therapeutically effective amount of a therapeutically effective amount of a first vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof, a therapeutically effective amount of a second vasodilating agent or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier. The classes and types of compounds which can be used in the method are described in the fourth aspect, above.

DETAILED DESCRIPTION


Humans include, of course, males and females. Although the pharmaceutical compositions of the present invention are envisaged primarily for the treatment of erectile dysfunction or male sexual dysfunction, they may also be useful for the treatment of female sexual dysfunction. Such female sexual dysfunction may include orgasmic dysfunction due to clitoral irregularities or disturbances.

Phentolamine, 3-[[[(4,5-dihydro-1H-imidazol-2-yl)methyl](4-methylphenyl)amino]phenol, and pharmaceutically acceptable salts, solvates, hydrates, crystalline polymorph forms and the free base thereof,

are useful in the treatment of sexual dysfunction. A rapidly disintegrating tablet and method of use to treat sexual dysfunction is disclosed in United States Patent No. 5,731,339, also incorporated herein by reference. Representative formulations comprising phentolamine are disclosed in U.S. 5,731,339. Phentolamine can exist in unsolvated as well as solvated forms, including hydrated forms, e.g. hemi-hydrate. In general, the solvated forms, with pharmaceutically acceptable solvents such as water, ethanol and the like are equivalent to the unsolvated forms for purposes of the invention. Phentolamine can form pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for salt formation are hydrohalic acids such as hydrochloric and hydrobromic; as well as other acids such as sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic, toluenesulfonic and other mineral and carboxylic acids known to those skilled in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to their respective free base form for purposes of this invention. Phentolamine can also form crystalline polymorph forms or crystalline forms thereof using suitable or conventional crystallization procedures.

The present invention is directed to the use of cyclic guanosine 3',5'-monophosphate phosphodiesterase (cGMP PDE) inhibitors in combination with the salts or esters of phentolamine, preferably, with phentolamine mesylate for the treatment of human sexual dysfunction, preferably erectial dysfunction. Examples of cGMP PDE inhibitors contemplated in this invention are as follows and are described in the following documents, as indicated. The disclosure of each of the below-referred to document is incorporated herein by reference.

European published application number 0201188, which discloses compounds of the formula




and the pharmaceutically acceptable salts thereof, in which:

R₁ is a lower alkyl of from one to six carbon atoms, a lower alkenyl of from one to six carbon atoms, a lower hydroxyalkyl of from one to six carbon atoms, a lower hydroxyalkenyl of from two to six carbon atoms, a lower aminoalkyl of from one to six carbon atoms, or a lower aminoalkenyl of from two to six carbon atoms;

n is 0 or an integer of from 1 to 4; and

Ar is a radical of the following general formula (R₂)



or 2, 3, or 4-pyridyl, in which X, Y, and Z are, independently, (1) hydrogen; (2) lower alkyl of from one to six carbon atoms; (3) halogen, (4) hydroxyl; (5) lower alkoxy of from one to six carbon atoms; - (6) nitro; (7) amino; (8) NR'R'' wherein R' and R'' are each, independently, (a) hydrogen or (b) lower alkyl of from one to six carbon atoms optionally substituted by (i) amino, (ii) morpholino or (iii) cycloalkyl of from, five to seven carbon atoms; (9) sulfonyl; or

(10) -SO₂NR'R'' wherein R' and R'' are as defined above;

with the proviso that not all of X, Y, and Z can be nitro, amino, or NR'R'' at once.

Preferred compounds include:

1-ethyl-3-methyl-5-phenylpyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-phenylpyrazolo[4,3-d]pyrimidine-7-
one;

1,3-dimethyl-5-(4-chlorophenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(4-methylphenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(4-nitrophenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(4-trifluoromethylphenyl)pyrazolo-
[4,3-d]-pyrimidine;

1,3-dimethyl-5-(4-aminophenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(3-aminophenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(3-nitrophenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(2-methoxyphenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(3,4-dichlorophenyl)pyrazolo[4,3-d]-
pyrimidine-7-one;

1,3-dimethyl-5-(3,4-dimethoxyphenyl)pyrazolo[4,3-
d]-pyrimidine-7-one;

1,3-dimethyl-5-(2,4-dimethoxyphenyl)pyrazolo[4,3-
d]-pyrimidine-7-one;

1,3-dimethyl-5-(2-nitro-4-chlorophenyl)pyrazolo-
[4,3-d]-pyrimidine-7-one;

1,3-dimethyl-5-(2-amino-4-chlorophenyl)pyrazolo-
[4,3-d]-pyrimidine-7-one;


1,3-dimethyl-5-(4-sulfonic acid phenyl)pyrazolo-
[4,3-d]-pyrimidine-7-one;

1,3-dimethyl-5-[4-(N-2-(dimethylamino)ethyl)-
benzenesulfonamide]pyrazolo[4,3-d]pyrimidine-7-
one;

1,3-dimethyl-5-(3,5-dimethoxyphenyl)pyrazolo[4,3-
d]-pyrimidine-7-one; or





1,3-dimethyl-5-(3-methoxyphenyl)pyrazolo[4,3-d]-
pyrimidine-7-one.

European published application number 0214708, which discloses compounds of the formula




in which:

A represents a group of formula:



or (e)



R¹ and R² are the same or different and each represents a hydrogen atom, a halogen atom or a group of formula -OR³;

R³ and R⁴ are the same or different and each represents a carbamoyl group or a carboxy group;

R⁵ and R⁶ both represent hydrogen atoms or together they represent an extra carbon-carbon bond between the carbon atoms to which they are attached;

R' represents a hydrogen atom, a halogen atom or a group of formula -OR', -NR''R''' or -SR';

R'' represents a halogen atom or a group of formula -OR'', -NR'''R'''' or -SR'';

R''' represents a hydrogen atom, a C₁-C₆ alkyl group, an alkylsulphonyl group, a haloalkylsulphonyl group, an arylsulphonyl group or a hydroxy-protecting group;

R'''' and R''''' are the same or different and each

represents a hydrogen atom, a hydroxy group, a C₁-C₆ alkyl group, a C₁-C₆ hydroxyalkyl group, a C₁-C₆ aminoalkyl group, an aralkyl group, an aryl group, a C₁-C₆ alkoxy group, an aralkyloxy group, an amino group, a C₁-C₆ aliphatic acyl group or an aromatic acyl group; or R'''' and R''''' together represent a substituted methylene group, or R'''' and R''''', together with the nitrogen atom to which they are attached, represent a heterocyclic group having 5 or 6 ring atoms, of which, in addition to the nitrogen atom shown, 0 or 1 are additional oxygen, nitrogen or sulphur hetero-atoms, said heterocyclic group being unsubstituted or having from 1 to 3 C₁-C₆ alkyl and/or C₁-C₆ alkoxy substituents;

R'''' represents a C₁-C₆ alkyl group;

Z represents a hydrogen atom, a hydroxy group or a substituted hydroxy group; and

W represents an alkoxy group or an aralkoxy group;

provided that, when A represents said group of

formula (e), R' and R'' both represent hydrogen atoms;

and pharmaceutically acceptable salts and esters thereof.

Preferred compounds include:

2-Amino-6-desamino-6-hydroxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-6-desamino-6-hydroxygriseolic acid 7'-amide and pharmaceutically acceptable salts and esters thereof.

2-Aminogriseolic acid and pharmaceutically acceptable salts and esters thereof.

Bis(pivaloyloxymethyl) 2-amino-6-desamino-6-hydroxygriseolate and pharmaceutically acceptable salts thereof.

2-Amino-N⁴-methoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-N⁴-benzyloxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Fluorogriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Chlorogriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-6-desamino-6-hydroxy-7'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-7'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Chloro-7'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-8-desamino-6-hydroxy-2'-chloro-2'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-6-desamino-6-hydroxy-2'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-2'-chloro-2'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-2'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Chloro-2'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

Griseolic acid N¹-oxide and pharmaceutically acceptable salts thereof.

2-Acetylamino-6-desamino-6-hydroxy-4',5'-dihydrogriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-6-desamino-6-hydroxy-4',5'-dihydrogriseolic acid and pharmaceutically acceptable salts and esters thereof.


2-Acetylamino-6-desamino-6-hydroxy-4',5'-dihydro-7'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2-Amino-6-desamino-6-hydroxy-4',5'-dihydro-7'-desoxygriseolic acid and pharmaceutically acceptable salts and esters thereof.

2,6-Dichloro-6-desamino-4',5'-dihydrogriseolic acid and pharmaceutically acceptable salts and esters thereof.


2-Chloro-4',5'-dihydrogriseolic acid and pharmaceutically acceptable salts and esters thereof.

European published application number 0319050, which discloses compounds of the formula



in which:


A represents a group of formula:



R^1 and R^2 are the same or different and each represents a hydrogen atom, a halogen atom or a group of formula $-OR^3$;
 R^3 and R^4 are the same or different and each represents a carbamoyl group or a carboxy group;
 R^5 and R^6 both represent hydrogen atoms;
 R^3 represents a hydrogen atom, a C_1-C_6 alkyl group, an alkylsulphonyl group, a haloalkylsulphonyl group, an arylsulphonyl group or a hydroxy-protecting group;
 R^{12} represents a C_1-C_6 alkyl group;
 and pharmaceutically acceptable salts and esters thereof.

-11-

European published application number 0293063, which discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein R¹ is C₁₋₆alkyl or C₂₋₆alkenyl, and R² is hydrogen or hydroxy.

Preferred compounds include:

2-(2-propoxyphenyl)-6-purinone,
2-(2-ethoxyphenyl)-6-purinone,
2-(2-butoxyphenyl)-6-purinone,
2-(2-isobutoxyphenyl)-6-purinone,
2-(2-propoxyphenyl)purine-6,8-dione,
2-(2-methoxyphenyl)purine-6,8-dione,
2-(2-ethoxyphenyl)purine-6,8-dione,
2-(2-butoxyphenyl)purine-6,8-dione,
2-(2-isobutoxyphenyl)purine-6,8-dione, or
2-(2-allyloxyphenyl)purine-6,8-dione
or a pharmaceutically acceptable salt thereof.

European published application number 0347027, which discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein
X is O or S;
R¹ is C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₆cycloalkyl, C₁₋₆alkyl, or C₁₋₆alkyl substituted by 1 to 6 fluoro groups;
R² is hydrogen, -CN, -CONR⁵R⁶, -CO₂R⁷, 5-tetrazolyl, -NO₂, -NH₂ or -NHCOR⁸ wherein R⁵, R⁶, R⁷ and R⁸ are independently hydrogen or C₁₋₆alkyl;
R³ is hydrogen or C₁₋₆alkyl; and
R⁴ is hydrogen or C₁₋₆alkyl;
with the proviso that R¹ is not methyl when R² is -CO₂H, -CO₂CH₂CH₃ or -CN, X is O, R³ is hydrogen and R⁴ is hydrogen or methyl.

-12-

Preferred compounds include:

3-cyano-6-(2-propoxyphenyl)-2(1H)-pyridinone,
 6-(2-propoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide,
 6-(2-propoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxylic acid,
 methyl 6-(2-propoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxylate,
 6-(2-propoxyphenyl)-3-(1H-tetrazol-5-yl)-2(1H)-pyridinone,
 6-(2-propoxyphenyl)-2(1H)-pyridinone,
 3-nitro-6-(2-propoxyphenyl)-2(1H)-pyridinone,
 3-cyano-6-(2-ethoxyphenyl)-2(1H)-pyridinone,
 3-amino-6-(2-propoxyphenyl)-2(1H)-pyridinone,
 3-cyano-4-methyl-6-(2-propoxyphenyl)-2(1H)-pyridinone,
 3-cyano-5-methyl-6-(2-propoxyphenyl)-2(1H)-pyridinone,
 3-cyano-6-(2-(1,1,2,3,3,3-hexafluoropropoxy)phenyl)-2(1H)-pyridinone,
 3-cyano-6-(2-propoxyphenyl)-2(1H)-pyridinethione,
 1,2-dihydro-4-methyl-2-oxo-6-(2-propoxyphenyl)pyridine-3-carboxylic acid,
 methyl 1,2-dihydro-4-methyl-2-oxo-6-(2-propoxyphenyl)pyridine-3-carboxylate,
 1,2-dihydro-4-methyl-2-oxo-6-(2-propoxyphenyl)pyridine-3-carboxamide,
 3-cyano-6-(2-cyclopropylmethoxyphenyl)-2(1H)-pyridinone,
 6-(2-butoxyphenyl)-3-cyano-2(1H)-pyridinone,
 6-(2-allyloxyphenyl)-3-cyano-2(1H)-pyridinone,
 3-cyano-6-[2-(2-methylpropoxy)phenyl]-2(1H)-pyridinone,
 6-(2-ethoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide,
 6-(2-cyclopropylmethoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide,
 6-(2-butoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide,
 6-(2-allyloxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide, or
 6-[2-(2-methylpropoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide,
 or a pharmaceutically acceptable salt thereof.

European published application number 0347146, which
 discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein




is a ring of sub-formula (a), (b), (c), (d), (e), (f) or (g) :


-13-




(a)




(b)




(c)




(d)



(e)



(f)




(g)

R¹ is C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₅cycloalkyl, C₁₋₆alkyl, or C₁₋₆alkyl substituted by 1 to 6 fluoro groups;
 R² is C₁₋₆alkylthio, C₁₋₆alkylsulphonyl, C₁₋₆alkoxy, hydroxy, hydrogen, hydrazino, C₁₋₆alkyl, phenyl, -NHCOR³ wherein R³ is hydrogen or C₁₋₆alkyl, or -NR⁴R⁵ wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, hexahydroazepino, morpholino or piperazino ring, or R⁴ and R⁵ are independently hydrogen, C₂₋₅cycloalkyl or C₁₋₆alkyl which is optionally substituted by -CF₃, phenyl, -S(O)_nC₁₋₆alkyl wherein n is 0, 1 or 2, -OR⁶, -CO₂R⁷ or -NR⁸R⁹ wherein R⁶ to R⁹ are independently hydrogen or C₁₋₆alkyl, provided that the carbon atom adjacent to the nitrogen atom is not substituted by said -S(O)_nC₁₋₆alkyl, -OR⁶ or -NR⁸R⁹ groups; and
 R is hydrogen and can also be hydroxy when R² is hydroxy.

Preferred compounds include:

2-(2-propoxyphenyl)pyrido[2,3-d]pyrimid-4(3H)-one,
 2-(2-propoxyphenyl)pyrido[3,4-d]pyrimid-4(3H)-one,
 2-(2-propoxyphenyl)pyrido[4,3-d]pyrimid-4(3H)-one,
 2-(2-propoxyphenyl)pyrido[3,2-d]pyrimid-4(3H)-one,
 2-(2-propoxyphenyl)pteridin-4(3H)-one,
 2-(2-propoxyphenyl)pteridin-4,6(3H,5H)-dione,
 2-(2-propoxyphenyl)pteridin-4,6,7(3H,5H,8H)-trione,
 5,6-dihydro-3-methylthio-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,
 3-amino-5,6-dihydro-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,
 3-methylamino-5,6-dihydro-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,
 3-methoxy-5,6-dihydro-5-oxo-7-(2-propoxyphenyl)pyrimido[5,4-e][1,2,4]triazine,
 3-methylthio-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,
 3-amino-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,
 3-methylamino-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,
 3-methoxy-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,
 3,8-dioxo-6-(2-propoxyphenyl)-3,4,7,8-tetrahydropyrimido[4,5-e][1,2,4]triazine,
 3-dimethylamino-8-oxo-6-(2-propoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,
 3-methylthio-8-oxo-6-(2-allyloxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,
 3-methylthio-8-oxo-6-(2-isobutoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine,
 3-methylthio-8-oxo-6-(2-cyclopropylmethoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine or
 3-methylthio-8-oxo-6-(2-methoxyphenyl)-7,8-dihydropyrimido[4,5-e][1,2,4]triazine
 or a pharmaceutically acceptable salt thereof.


European published application number 0349239, which discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein




is a ring of sub-formula (a), (b) or (c):



(a)



(b)



(c),


X is oxygen or sulphur, and

R¹ is C₁-₄alkyl, C₂-₆alkenyl, C₃-₅cycloalkyl, C₁-₄alkyl, or C₁-₄alkyl substituted by 1 to 8 fluoro groups.

Preferred compounds include:

6-(2-propoxyphenyl)pyrazolo[3,4-d]pyrimidin-4(5H)-one,
2-(2-propoxyphenyl)thieno[2,3-d]pyrimidin-4(3H)-one,
2-(2-propoxyphenyl)[1,2,5]oxadiazolo[3,4-d]pyrimidin-4(3H)-one, or
2-(2-propoxyphenyl)[1,2,5]thiadiazolo[3,4-d]pyrimidin-4(3H)-one,
or a pharmaceutically acceptable salt thereof.

European published application number 0351058, which discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein


R¹ is C₁-εalkyl, C₂-εalkenyl, C₂-εcycloalkyl, C₁-εalkyl, or C₁-εalkyl substituted by 1 to 6 fluoro groups; R² is C₁-εalkylthio, C₁-εalkylsulphonyl, C₁-εalkoxy, hydroxy, hydrogen, hydrazino, C₁-εalkyl, phenyl, -NHCOR³ wherein R³ is hydrogen or C₁-εalkyl, or -NR⁴R⁵, wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, hexahydroazepino, morpholino or piperazino ring, or R⁴ and R⁵ are independently hydrogen, C₂-εcycloalkyl or C₁-εalkyl which is optionally substituted by -CF₃, phenyl, -S(O)ₙC₁-εalkyl wherein n is 0, 1 or 2, -OR⁶, -CO₂R⁷ or -NR⁸R⁹ wherein R⁶ to R⁹ are independently hydrogen or C₁-εalkyl, provided that the carbon atom adjacent to the nitrogen atom is not substituted by said -S(O)ₙC₁-εalkyl, -OR⁶ or -NR⁸R⁹ groups; and



is a ring of sub-formula (a) or (b) :



(a)




(b) .

Preferred compounds include:

7-methylthio-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methylthio-2-(2-ethoxyphenyl)-4-oxo-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methylthio-2-(2-methoxyphenyl)-4-oxo-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methylthio-2-(2-isobutoxyphenyl)-4-oxo-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methylthio-2-(2-cyclopropylmethoxyphenyl)-4-oxo-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methylthio-2-(2-allyloxyphenyl)-4-oxo-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-amino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methylamino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-dimethylamino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-hydrazino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-ethylamino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-hydroxyethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-ethyl-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methylamino-2-(2-methoxyphenyl)-4-oxo-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-phenyl-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,

7-morpholino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-cyclopropylamino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-acetamido-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-propylamino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(3-hydroxypropylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-methoxyethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-dimethylaminoethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-hydroxypropylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(3-methylthiopropylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-aminoethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine hydrochloride,
 7-(3-methylsulfinylpropylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(3-methylsulphonylpropylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 4,7-dioxo-2-(2-propoxyphenyl)-3,4,7,8-tetrahydropyrimido[4,5-d]pyrimidine,
 7-methylsulphonyl-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-diethylamino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-ethoxycarbonyl-ethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(ethoxycarbonylmethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-carboxyethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(carboxymethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-ethoxy-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-methoxy-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2,2,2-trifluoroethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-propoxy-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(N-ethyl-N-hydroxyethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-dipropylamino-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine,
 7-(2-phenethylamino)-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidine, or
 4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido[5,4-d]pyrimidine,
 or a pharmaceutically acceptable salt thereof.

European published application number 0352960, which discloses compounds of the formula



or a pharmaceutically acceptable salt thereof, wherein

R¹ is C₁-alkyl, C₂-alkenyl, C₃-cycloalkyl, C₁-alkyl, phenyl, C₁-alkyl or C₁-alkyl substituted by 1 to 6 fluoro groups;

R² is hydrogen, hydroxy, C₁-alkyl, phenyl, mercapto, C₁-alkylthio, CF₃ or amino;

R³ is hydrogen, nitro, amino, C₁-alkanoylamino, C₁-alkoxy, C₁-alkyl, halo, SO₂NR⁴R⁵, CONR⁴R⁵,

cyano or C₁-alkylS(O)_n;

R⁴ and R⁵ are independently hydrogen or C₁-alkyl; and


n is 0, 1 or 2;

provided that R³ is not hydrogen when R¹ is C₁-alkyl or C₂-alkenyl and R² is hydrogen or hydroxy.

Preferred compounds include:

2-(2-{2,2,2-trifluoroethoxy}phenyl)purin-6-one,
 2-(2-cyclopropylmethoxyphenyl)purin-6-one,
 2-(2-cyclopropylmethoxyphenyl)purin-8,8-dione,
 2-(2-benzyloxyphenyl)purin-6,8-dione,
 2-(2-propoxyphenyl)-8-trifluoromethylpurin-8-one,
 2-(2-propoxyphenyl)-8-phenylpurin-6-one,
 2-(2-propoxyphenyl)-8-methylpurin-6-one,
 2-(2-propoxyphenyl)-8-mercaptopurin-6-one,
 2-(2-propoxyphenyl)-8-methylthiopurin-6-one,
 2-(2-propoxyphenyl)-8-aminopurin-6-one,
 2-(2-propoxy-5-nitrophenyl)purin-6-one,
 2-(2-propoxy-5-aminophenyl)purin-6-one,
 2-(2-propoxy-5-acetamidophenyl)purin-6-one,
 2-(2-propoxy-4-methoxyphenyl)purin-8-one,
 2-(2-propoxy-5-methoxyphenyl)purin-8-one,
 2-(2-propoxy-5-chlorophenyl)purin-8-one,
 2-(2-propoxy-4-methylphenyl)purin-6-one,
 2-(2-propoxy-5-fluorophenyl)purin-8-one,
 2-(2-propoxy-5-dimethylsulphamoylphenyl)purin-6-one,
 2-(2-propoxy-5-methylsulphamoylphenyl)purin-6-one,
 2-(2-propoxy-5-sulphamoylphenyl)purin-8-one,
 2-(2-propoxy-4-methylthiophenyl)purin-6-one,
 2-(2-propoxy-5-cyanophenyl)purin-8-one, or
 2-(2-propoxy-5-carbamoylphenyl)purin-6-one,
 or a pharmaceutically acceptable salt thereof.

European published application number 0371731, which discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein
 R¹ is C₁-alkyl, C₂-alkenyl, C₃-cycloalkyl, C₁-alkyl, phenyl, C₁-alkyl or C₁-alkyl substituted by 1 to 6 fluoro groups;
 R² is hydrogen, C₁-alkyl, C₁-alkylthio, C₁-alkoxy, nitro or -NR³R⁴; and
 R³ and R⁴ are independently hydrogen or C₁-alkyl optionally substituted by hydroxy provided that the carbon atom adjacent to the nitrogen atom is not substituted by hydroxy;
 with the proviso that R¹ is not methyl or ethyl when R² is hydrogen.

Preferred compounds include:

2-(2-propoxyphenyl)quinazolin-4(3H)-one,
 7-methylthio-2-(2-propoxyphenyl)quinazolin-4(3H)-one,
 7-nitro-2-(2-propoxyphenyl)-4(3H)-quinazolinone,
 7-amino-2-(2-propoxyphenyl)-4(3H)-quinazolinone, or
 7-methylamino-2-(2-propoxyphenyl)-4(3H)-quinazolinone
 or a pharmaceutically acceptable salt thereof.

European published application number 0395328, which
 discloses compounds of the formula



or a pharmaceutically acceptable salt thereof, wherein


R¹ is C₁-₆alkyl, C₂-₆alkenyl, C₃-₆cycloalkyl, C₁-₆alkyl, phenyl, C₁-₆alkyl or C₁-₆alkyl substituted by 1 to 6 fluoro groups; and

R² is C₁-₆alkyl, phenyl, hydroxy, C₁-₆alkoxy, halo, -NHCOR³, -NHCONHR⁴, 5-tetrazolyl, -CO₂R⁵, cyano, -CONR⁶R⁷, or -NR⁸R⁹ wherein R³ to R⁹ are independently hydrogen or C₁-₆alkyl and R⁸ and R⁹ are independently hydrogen or C₁-₆alkyl optionally substituted by hydroxy provided that the carbon atom adjacent to the nitrogen atom is not substituted by hydroxy;

Preferred compounds include:

6-amino-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 6-acetamido-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 6-propionamido-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 6-butyramido-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 6-N-methylureido-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 4,6-dihydroxy-2-(2-propoxyphenyl)pyrimidine,
 4-chloro-6-hydroxy-2-(2-propoxyphenyl)pyrimidine,
 6-ethylamino-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 6-propylamino-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 6-(2-hydroxyethylamino)-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 6-(3-hydroxypropylamino)-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 4-hydroxy-6-methyl-2-(2-propoxyphenyl)pyrimidine,
 6-hydroxy-2-(2-propoxyphenyl)pyrimidine-4-carboxylic acid,
 ethyl 6-hydroxy-2-(2-propoxyphenyl)pyrimidine-4-carboxylate,
 6-hydroxy-2-(2-propoxyphenyl)pyrimidine-4-carboxamide,
 4-cyano-6-hydroxy-2-(2-propoxyphenyl)pyrimidine,
 2-(2-propoxyphenyl)-6-(1H-tetrazol-5-yl)pyrimidin-4(3H)-one,
 4-ethyl-6-hydroxy-2-(2-propoxyphenyl)pyrimidine,
 4-hydroxy-6-phenyl-2-(2-propoxyphenyl)pyrimidine,
 N-methyl 6-hydroxy-2-(2-propoxyphenyl)pyrimidine-4-carboxamide,
 N-ethyl 6-hydroxy-2-(2-propoxyphenyl)pyrimidine-4-carboxamide,
 N-propyl 6-hydroxy-2-(2-propoxyphenyl)pyrimidine-4-carboxamide,
 6-ethoxy-2-(2-propoxyphenyl)pyrimidin-4(3H)-one, or
 6-N,N-bis-(2-hydroxyethyl)amino-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 or a pharmaceutically acceptable salt thereof.

European published application number 0400583, which discloses compounds of the formula




wherein -

A is N or CH;



B is N CR₃;

D is N or CR₂;

R, R₁, are the same or independently hydrogen, hydroxy, loweralkyl, lower alkoxy, phenyloxy, R_cS(O)_n-, W-ALK-Q-,



R₂ is hydrogen, lower alkyl, phenyl which may be substituted by up to three methoxy groups, lower alkyl substituted by phenyl which may be substituted by up to three methoxy groups, - lower alkyl -N(R₈)₂,




pyridinyl or lower-alkyl pyridinyl;

R₃ is hydrogen, lower alkyl, phenyl, lower alkylphenyl, pyridinyl or loweralkyl pyridinyl;

R₄, R₅ are the same or independently hydrogen or lower alkyl;


R₆ is lower alkyl, phenyl, lower alkylphenyl or pyridinyl;

R₇ are the same or independently hydrogen, loweralkyl, phenyl, pyridinyl,




R₈ are the same or independently lower alkyl, phenyl or pyridinyl;

-20-



W is hydroxy, loweralkoxy, phenoxy, $-N(R_{10})_2$, $-N$ (cyclopentyl), $-N$ (cyclohexyl)X,



ALK is a C_1 - C_4 straight or branched chain alkyl;

R_3 is hydrogen, lower alkyl or phenyl;

R_{10} are the same or independently hydrogen, loweralkyl or phenyl;

R_{11} are the same or independently hydrogen or lower alkyl;

X is $-CH_2-$, $-O-$, $S(O)_n$, $-NR_{10}$;


n is the integer 0, 1 or 2 and

p is the integer 0 or 1.

with the provisos that:

a) one and only one of B or D must be N;

b) when A is CH, when D is N, when B is CR_3 where R_3 is H, when R_2 is hydrogen, lower alkyl or phenyl then R and/or R_1 must be



or W-ALK-Q;


and the pharmaceutically acceptable salts thereof.

Preferred compounds include:

1-ethyl-8-(1H-imidazol-1-yl)-3-methylimidazo[1,5-a]quinoxalin-4(5H)-one, 1-ethyl-8-(1H-imidazol-1-yl)imidazo[1,5-a]quinoxalin-4(5H)-one, 1-ethyl-3-methyl-8-(4-morpholino)imidazo[1,5-a]quinoxalin-4(5H)-one, 1-ethyl-8-(2-ethyl-4-methyl-1H-imidazol-1-yl)-3-methylimidazo[1,5-a]quinoxalin-4(5H)-one, 1-methyl-8-(2-methyl-1H-imidazol-1-yl)imidazo[1,5-a]quinoxalin-4(5H)-one, 8-(1H-imidazol-1-yl)-1-methylimidazo[1,5-a]quinoxalin-4(5H)-one, 1-ethyl-3-methyl-8-(pyrrolidin-1-yl)imidazo[1,5-a]quinoxalin-4(5H)-one, 1-((morpholin-4-yl)methyl)imidazo[1,5-a]quinoxalin-4(5H)-one, or 6-ethoxy-1-ethyl-8-(2-ethyl-4-methyl-1H-imidazol-1-yl)-3-methylimidazo[1,5-a]quinoxalin-4(5H)-one, 8-(1H-imidazol-1-yl)imidazo[1,2-a]quinoxalin-4(5H)-one, 8-(1H-imidazol-1-yl)imidazo[1,2-a]quinoxalin-4(5H)-one, 8-ethylimidazo[1,5-a]pyrido[3,2-e]pyrazin-6(5H)-one, 8-methyl-2-(2-methyl-1H-imidazol-1-yl)imidazo[1,5-a]pyrido[3,2-e]pyrazin-6(5H)-one, 8-((2-ethyl-1H-imidazol-1-yl)methyl)imidazo[1,5-a]pyrido[3,2-e]pyrazin-6(5H)-one, or 1-ethylimidazo[1,5-a]pyrido[4,3-e]pyrazin-4(5H)-one, 1-ethylimidazo[1,2-a]pyrido[3,2-e]pyrazin-6(5H)-one, 2-phenylimidazo[1,2-a]pyrido[2,3-e]pyrazin-4(5H)-one, or 2-(1H-imidazol-1-yl)imidazo[1,2-a]pyrido[3,2-e]pyrazin-6(5H)-one.

-21-

European published application number 0400799, which discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein R¹ is C₁-₆ alkyl, C₂-₆ alkenyl, C₃-₅ cycloalkyl, C₁-₆ alkyl, phenyl, C₁-₆ alkyl or C₁-₆ alkyl substituted by 1 to 6 fluoro groups; and R² is hydrogen, amino, -NHCOR³, or -CONR⁴R⁵, wherein R³ is C₁-₆ alkyl, R⁴ is C₁-₆ alkyl and R⁵ is hydrogen or C₁-₆ alkyl.

Preferred compounds include:

1,6-dihydro-6-oxo-2-(2-propoxyphenyl)pyrimidine-5-carboxamide,
 N-methyl 1,6-dihydro-6-oxo-2-(2-propoxyphenyl)pyrimidine-5-carboxamide,
 N,N-dimethyl 1,6-dihydro-6-oxo-2-(2-propoxyphenyl)pyrimidine-5-carboxamide,
 5-amino-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 5-acetamido-2-(2-propoxyphenyl)pyrimidin-4(3H)-one, or
 2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
 or a pharmaceutically acceptable salt thereof.

European published application number 0428268, which discloses compounds of the formula



or a pharmaceutically acceptable salt thereof, wherein
 X is O or S;
 R¹ is C₁-6 alkyl, C₂-6 alkenyl, C₃-5 cycloalkyl, C₁-4 alkyl, or C₁-4 alkyl substituted by 1 to 3 fluoro groups;
 R² is hydrogen, -CN, -CONR⁵R⁶, -CO₂R⁷, 5-tetrazolyl, -NO₂, -NH₂ or -NHCOR⁸ wherein R⁵ to R⁸ are independently hydrogen or C₁-4 alkyl;
 R³ is hydrogen or C₁-4 alkyl;
 R⁴ is hydrogen or C₁-4 alkyl; and
 R is halo, C₁-4 alkyl, C₁-4 alkoxy, cyano, -CONR⁹R¹⁰, -CO₂R¹¹, -S(O)_nC₁-4 alkyl, -NO₂, -NH₂, -NHCOR¹², or -SO₂NR¹³R¹⁴ wherein n is 0, 1 or 2 and R⁹ to R¹⁴ are independently hydrogen or C₁-4 alkyl; with the proviso that R¹ is not methyl when R² is -CO₂H, -CO₂CH₂CH₃ or -CN, X is O, R³ is hydrogen, R⁴ is hydrogen or methyl and R is 6-methoxy.


Preferred compounds include:

3-cyano-6-(2-methoxy-4-methylthiophenyl)-2(1H)-pyridinone,
 3-cyano-6-(4-methylthio-2-propoxyphenyl)-2(1H)-pyridinone,
 1,2-dihydro-6-(4-methylthio-2-propoxyphenyl)-2-oxo-3-pyridine carboxamide,
 3-cyano-6-(2-methoxy-4-methylsulphonylphenyl)-2(1H)-pyridinone,
 3-cyano-6-(4-methylsulphonyl-2-propoxyphenyl)-2(1H)-pyridinone,
 3-cyano-6-(4-methylsulphonyl-2-propoxyphenyl)-2(1H)-pyridinone,
 3-cyano-6-(2-methoxy-4-methylsulphonylphenyl)-2(1H)-pyridinone,
 3-cyano-6-(5-fluoro-2-propoxyphenyl)-2(1H)-pyridinone,
 1,2-dihydro-6-(5-fluoro-2-propoxyphenyl)-2-oxo-3-pyridine carboxamide,
 3-cyano-6-(4-methoxy-2-propoxyphenyl)-2(1H)-pyridinone,
 1,2-dihydro-6-(4-methoxy-2-propoxyphenyl)-2-oxo-3-pyridine carboxamide,
 3-cyano-6-(5-methoxy-2-propoxyphenyl)-2(1H)-pyridinone,
 1,2-dihydro-6-(5-methoxy-2-propoxyphenyl)-2-oxo-3-pyridine carboxamide,
 3-cyano-6-(5-cyano-2-propoxyphenyl)-2(1H)-pyridinone,
 3-(3-carboxamido-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxybenzamide,
 methyl 3-(3-cyano-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxybenzoate,
 3-(3-cyano-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxybenzamide,
 N-methyl 3-(3-cyano-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxybenzamide,
 N-methyl 3-(3-carboxamido-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxybenzamide,
 N,N-dimethyl 3-(3-cyano-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxybenzamide,
 N,N-dimethyl 3-(3-carboxamido-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxybenzamide,
 4-(3-cyano-1,2-dihydro-2-oxo-6-pyridinyl)-3-propoxybenzonitrile,
 4-(3-carboxamido-1,2-dihydro-2-oxo-6-pyridinyl)-3-propoxybenzamide,

-23-

3-cyano-6-(5-methylthio-2-propoxyphenyl)-2(1H)pyridinone,
 3-(3-cyano-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxy-N,N-dimethylbenzenesulphonamide,
 3-(3-carboxamido-1,2-dihydro-2-oxo-6-pyridinyl)-4-propoxy-N,N-dimethylbenzenesulphonamide,
 6-(2-cyclopropylmethoxy-5-fluorophenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide,
 6-(5-fluoro-2-(2-methylpropoxy)phenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide,
 3-cyano-6-(5-nitro-2-propoxyphenyl)-2(1H)-pyridinone,
 1,2-dihydro-6-(5-nitro-2-propoxyphenyl)-2-oxo-3-pyridinone carboxamide,
 3-cyano-6-(5-amino-2-propoxyphenyl)-2(1H)-pyridinone,
 1,2-dihydro-6-(5-amino-2-propoxyphenyl)-2-oxo-3-pyridinone carboxamide,
 3-cyano-6-(5-acetamido-2-propoxyphenyl)-2(1H)-pyridinone or
 1,2-dihydro-6-(5-acetamido-2-propoxyphenyl)-2-oxo-3-pyridine carboxamide,
 or a pharmaceutically acceptable salt thereof.

European published application number 0442204, which
 discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein

R¹ is C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆cycloalkyl, C₁₋₆alkyl, or C₁₋₆alkyl substituted by 1 to 6 fluoro groups;
 R² is C₁₋₆alkylthio, C₁₋₆alkylsulphonyl, C₁₋₆alkoxy, hydroxy, hydrogen, hydrazino, C₁₋₆alkyl, phenyl, -NHCOR³ wherein R³ is hydrogen or C₁₋₆alkyl, or -NR⁴R⁵, wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, hexahydroazepino, morpholino or piperazino ring, or R⁴ and R⁵ are independently hydrogen, C₂₋₆cycloalkyl or C₁₋₆alkyl which is optionally substituted by -CF₃, phenyl, -S(O)_nC₁₋₆alkyl wherein


n is 0, 1 or 2, -OR⁶, -CO₂R⁷ or -NR⁸R⁹ wherein R⁶ to R⁹ are independently hydrogen or C₁₋₆alkyl, provided that the carbon atom adjacent to the nitrogen atom is not substituted by said -S(O)_nC₁₋₆alkyl, -OR⁶ or -NR⁸R⁹ groups;

R is halo, C₁₋₆alkyl, C₁₋₆alkoxy, cyano, -CONR¹⁰R¹¹, CO₂R¹², C₁₋₆alkylS(O)_n, -NO₂, -NH₂, -NHCOR¹³ or SO₂NR¹⁴R¹⁵ wherein n is 0, 1 or 2 and R¹⁰ to R¹⁵ are independently hydrogen or C₁₋₆alkyl; and

A is a ring of sub-formula (a) or (b) :




(a)



(b)

European published application number 0579496, which discloses compounds of the formula



wherein — represents a single or double bond;

R^1 is hydrogen or C_{1-4} alkyl;

Y is a single bond or C_{1-6} alkylene;

A is

- (i) $\text{—CyA—}(R^2)_1$,
- (ii) —O—R^0 or $\text{—S(O)}_p\text{—R}^0$, or
- (iii) $\text{—NR}^{16}\text{R}^{17}$;

in which R^0 is hydrogen, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl or $\text{—CyA—}(R^2)_1$;

R^{16} and R^{17} independently are hydrogen or C_{1-4} alkyl;

p is 0-2;

CyA is

- (1) a 3-7 membered, saturated or unsaturated carbocycle,
- (2) a 4-7 membered, unsaturated or partially saturated heterocycle containing one nitrogen atom,
- (3) a 4-7 membered, unsaturated or partially saturated heterocycle containing one nitrogen atom and one oxygen atom,
- (4) a 4-7 membered, unsaturated or partially saturated heterocycle containing one nitrogen atom and two oxygen atoms,
- (5) a 4-7 membered, unsaturated or partially saturated heterocycle containing two nitrogen atoms and one oxygen atom,
- (6) a 4-7 membered, unsaturated or partially saturated heterocycle containing one or two sulfur atoms,
- (7) a 4-7 membered, unsaturated, partially saturated or fully saturated heterocycle containing one or two oxygen atoms;

R^2 is (1) hydrogen, (2) C_{1-4} alkyl, (3) C_{1-4} alkoxy, (4) —COOR^6 , in which R^6 is hydrogen or C_{1-4} alkyl, (5) $\text{—NR}^6\text{R}^7$, in which R^6 and R^7 independently are hydrogen or C_{1-4} alkyl, (6) $\text{—SO}_2\text{NR}^6\text{R}^7$, in which R^6 and R^7 are as hereinbefore defined, (7) halogen, (8) trifluoromethyl, (9) nitro or (10) trifluoromethoxy;

Z is a single bond, methylene, ethylene, vinylene or ethynylene;

CyB is

- (1) a 4-7 membered, unsaturated or partially saturated heterocycle containing one nitrogen atom,
- (2) a 4-7 membered, unsaturated or partially saturated heterocycle containing two nitrogen atoms,
- (3) a 4-7 membered, unsaturated or partially saturated heterocycle containing three nitrogen atoms,
- (4) a 4-7 membered, unsaturated or partially saturated heterocycle containing one or two oxygen atoms,
- (5) a 4-7 membered, unsaturated or partially saturated heterocycle containing one or two sulfur atoms,

R^3 is hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, halogen or trifluoromethyl;

R^4 is (1) hydrogen, (2) C_{1-4} alkyl, (3) C_{1-4} alkoxy, (4) —COOR^8 , in which R^8 is hydrogen or C_{1-4} alkyl, (5) $\text{—NR}^8\text{R}^{10}$, in which R^8 is hydrogen, C_{1-4} alkyl or phenyl(C_{1-4} alkyl) and R^{10} is hydrogen or C_{1-4} alkyl, (6) —NHCOR^{11} , in which R^{11} is C_{1-4} alkyl, (7) $\text{—NH—SO}_2\text{R}^{11}$, in which R^{11} is as hereinbefore defined, (8) $\text{SO}_2\text{NR}^9\text{R}^{10}$ in which R^9 and R^{10} are as hereinbefore defined, (9) —OCOR^{11} , in which R^{11} is as hereinbefore defined, (10) halogen, (11) trifluoromethyl, (12) hydroxy, (13) nitro, (14) cyano, (15) $\text{—SO}_2\text{N=CHNR}^{12}\text{R}^{13}$ in which R^{12} is hydrogen or C_{1-4} alkyl and R^{13} is C_{1-4} alkyl, (16) $\text{—CONR}^{14}\text{R}^{15}$ in which R^{14} is hydrogen or C_{1-4} alkyl or phenyl(C_{1-4} alkyl) and R^{15} is C_{1-4} alkyl or (17) C_{1-4} alkylthio, (18) C_{1-4} alkylsulfinyl, (19) C_{1-4} alkylsulfonyl, (20) ethynyl, (21) hydroxymethyl, (22) tri(C_{1-4} alkyl)silyl ethynyl or (23) acetyl;

and l , m and n independently are 1 or 2;

with the proviso that

- (1) $\text{CyA—}(R^2)_1$ does not represent cyclopentyl or trifluoromethylphenyl when Y is a single bond,
 - (2) CyB does not bond to Z through a nitrogen atom when Z is vinylene or ethynylene,
 - (3) CyB is not pyridine or thiophene when CyA is a 4-7 membered unsaturated, partially saturated or fully saturated heterocycle containing one or two oxygen atoms, and
 - (4) Y is not a single bond when A is (i) —O—R^0 or $\text{—S(O)}_p\text{—R}^0$ or (iii) $\text{—NR}^{16}\text{R}^{17}$;
- or a pharmaceutically acceptable salt thereof, or a hydrate thereof.

Preferred compounds include:


- 4-phenylmethylanino-2-(3-pyridyl)quinazoline,
- 4-(3-methylphenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(3,4-dimethoxyphenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(4-carboxyphenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(3-methoxycarbonylphenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(4-(N,N-dimethylamino)phenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(4-sulfamoylphenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(3-chlorophenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(3-trifluoromethylphenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-(3-nitrophenylmethyl)amino-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-2-(6-methyl-3-pyridyl)quinazoline,
- 4-phenylmethylanino-2-(6-methoxy-3-pyridyl)quinazoline,
- 4-phenylmethylanino-2-(6-chloro-3-pyridyl)quinazoline,
- 4-phenylmethylanino-2-(6-trifluoromethyl-3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methyl-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methoxy-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6,7-dimethoxy-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-carboxy-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methoxycarbonyl-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-amino-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-(N,N-dimethylamino)-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-acetylamino-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methanesulfonylamino-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-sulfamoyl-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-acetoxy-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-chloro-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-bromo-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-7-fluoro-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-trifluoromethyl-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-trifluoromethoxy-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-hydroxy-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-nitro-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-cyano-2-(3-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methyl-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methoxy-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6,7-dimethoxy-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-carboxy-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methoxycarbonyl-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-amino-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-(N,N-dimethylamino)-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-acetylamino-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-methanesulfonylamino-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-sulfamoyl-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-acetoxy-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-chloro-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-bromo-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-7-fluoro-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-trifluoromethyl-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-trifluoromethoxy-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-hydroxy-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-nitro-2-(4-pyridyl)quinazoline,
- 4-phenylmethylanino-6-cyano-2-(4-pyridyl)quinazoline,
- 4-phenylamino-2-(3-pyridyl)quinazoline,
- 4-(3-methoxycarbonylphenyl)amino-2-(3-pyridyl)quinazoline,
- 4-phenylethylanino-2-(3-pyridyl)quinazoline,

4-phenylmethylanino-2-(2-pyridyl)quinazoline,
4-phenylmethylanino-2-(4-pyridyl)quinazoline,
4-phenylmethylanino-2-(2-(3-pyridyl)ethyl)quinazoline,
4-phenylmethylanino-2-(2-(3-pyridyl)vinyl)quinazoline,
6-iodo-4-phenylmethylanino-2-(3-pyridyl)quinazoline,
4-(3-carboxyphenyl)amino-2-(4-pyridyl)quinazoline,
6-fluoro-4-phenylmethylanino-2-(3-pyridyl)quinazoline,
4-(cyclopropylmethyl)amino-2-(3-pyridyl)quinazoline,
4-(cyclohexylmethyl)amino-2-(3-pyridyl)quinazoline,
4-(2-azepinylmethyl)amino-2-(3-pyridyl)quinazoline,
4-(3-pyridylmethyl)amino-2-(3-pyridyl)quinazoline,
4-((1-methyl-2-pyrrolyl)methyl)amino-2-(3-pyridyl)quinazoline,
4-(3-isoxazolyl)amino-2-(3-pyridyl)quinazoline,
4-(3-isoxazolylmethyl)amino-2-(3-pyridyl)quinazoline,
4-(2-thienylmethyl)amino-2-(3-pyridyl)quinazoline,
4-(2-furylmethyl)amino-2-(1-imidazolyl)quinazoline,
4-(2-tetrahydrofuranlylmethyl)amino-2-(1-imidazolyl)quinazoline,
4-(4-tetrahydropyranlylmethyl)amino-2-(1-imidazolyl)quinazoline,
6-methoxy-4-(4-tetrahydropyranlylmethyl)amino-2-(1-imidazolyl)quinazoline,
6-chloro-4-(4-tetrahydropyranlylmethyl)amino-2-(1-imidazolyl)quinazoline,
4-(2-phenoxyethyl)amino-2-(1-imidazolyl)quinazoline,
4-(2-thienylethyl)amino-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-2-(1-imidazolyl)quinazoline,
4-(1,1-dimethyl-2-methoxyethyl)amino-2-(1-imidazolyl)quinazoline,
6-methoxy-4-(2-methoxyethyl)amino-2-(1-imidazolyl)quinazoline,
6-chloro-4-(2-methoxyethyl)amino-2-(1-imidazolyl)quinazoline,
4-(3-ethoxypropyl)amino-2-(1-imidazolyl)quinazoline,
6-nitro-4-(2-methoxyethyl)amino-2-(1-imidazolyl)quinazoline,
6-chloro-4-(2-ethoxyethyl)amino-2-(3-pyridyl)quinazoline,
6,7-dimethoxy-4-(2-methoxyethyl)amino-2-(1-imidazolyl)quinazoline,
6-chloro-4-(2-(2-hydroxyethoxy)ethyl)amino-2-(1-imidazolyl)quinazoline,
6-chloro-4-(2-dimethylaminoethyl)amino-2-(1-imidazolyl)quinazoline,
6-methoxy-4-(2-(2-hydroxyethoxy)ethyl)amino-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6-iodo-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6-methoxy-2-(2-methyl-1-imidazolyl)quinazoline,
4-(2-hydroxyethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6,8-diiodo-2-(1-imidazolyl)quinazoline,
4-(2-(2-hydroxyethoxy)ethyl)amino-6-iodo-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6-methylthio-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6-methylsulfinyl-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6-methylsulfonyl-2-(1-imidazolyl)quinazoline,
4-(2-(2-hydroxyethoxy)ethyl)amino-6-methylsulfinyl-2-(1-imidazolyl)quinazoline,
2-(1-imidazolyl)-4-(2-methoxyethyl)amino-6-(2-triethylsilylethynyl)quinazoline,
6-acetyl-4-(2-methoxyethyl)amino-2-(3-pyridyl)quinazoline,
6-ethynyl-4-(2-methoxyethyl)amino-2-(3-pyridyl)quinazoline,
4-[2-(2-hydroxyethoxy)ethyl]amino-6-acetyl-2-(1-imidazolyl)quinazoline,
4-(2-methylthioethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
4-(2-methylsulfinylethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
4-(2-methylsulfonylethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
4-[2-(2-hydroxyethoxy)ethyl]amino-6-methoxycarbonyl-2-(1-imidazolyl)quinazoline,
4-[2-(2-hydroxyethoxy)ethyl]amino-6-hydroxymethyl-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6-hydroxymethyl-2-(1-imidazolyl)quinazoline,
4-(2-methoxyethyl)amino-6-methoxycarbonyl-2-(1-imidazolyl)quinazoline,
4-(3-methoxypropyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
4-(2-(2-hydroxyethoxy)ethyl)amino-6-methylthio-2-(1-imidazolyl)quinazoline,
2-(1-imidazolyl)-4-[2-(2-hydroxyethoxy)ethyl]amino-6-(2-triisopropylsilylethynyl)quinazoline,
4-phenylmethylanino-6-methyl-2-(1-imidazolyl)quinazoline,
4-phenylmethylanino-6-methoxy-2-(1-imidazolyl)quinazoline,
4-phenylmethylanino-6,7-dimethoxy-2-(1-imidazolyl)quinazoline,
4-phenylmethylanino-6-carboxy-2-(1-imidazolyl)quinazoline,
4-phenylmethylanino-6-methoxycarbonyl-2-(1-imidazolyl)quinazoline,

4-phenylmethylamino-6-amino-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-(N,N-dimethylamino)-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-acetylamino-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-methanesulfonylamino-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-sulfamoyl-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-acetoxy-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-chloro-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-bromo-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-7-fluoro-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-trifluoromethyl-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-trifluoromethoxy-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-hydroxy-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-nitro-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6-cyano-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-2-((1-imidazolyl)methyl)quinazoline,
4-phenylmethylamino-2-(2-methyl-1-imidazolyl)quinazoline,
6-bromo-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
7-chloro-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-chloro-4-phenylamino-2-(1-imidazolylmethyl)quinazoline,
6-nitro-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-methoxy-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-chloro-4-phenylmethylamino-2-(1-imidazolylmethyl)quinazoline,
6-chloro-4-(3-carboxyphenyl)amino-2-(1-imidazolylmethyl)quinazoline,
6-dimethylaminosulfonyl-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6,7-dimethoxy-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
4-(3,4-dimethoxyphenylmethyl)amino-2-(1-imidazolyl)quinazoline,
6-dimethylaminomethylideneaminosulfonyl-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-(phenylmethylaminosulfonyl)-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
4-(2-phenylethyl)amino-2-(1-imidazolyl)quinazoline,
4-cyclohexylmethylamino-2-(1-imidazolyl)quinazoline,
6-carboxy-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-phenylmethylaminocarbonyl-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-iodo-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-ethoxycarbonyl-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
6-hydroxy-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
4-(4-trifluoromethoxyphenylmethyl)amino-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-2-(2-azepinyl)quinazoline,
4-phenylmethylamino-2-(1,5-diazepin-2-yl)quinazoline,
4-phenylmethylamino-2-(2-pyrimidinyl)quinazoline,
4-phenylmethylamino-2-(2-triazinyl)quinazoline,

4-phenylmethylamino-2-(2-pyrrolyl)quinazoline,
4-phenylmethylamino-2-(1-triazolyl)quinazoline,
6-hydroxy-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
4-(3-trifluoromethoxyphenylmethyl)amino-2-(1-imidazolyl)quinazoline,
4-phenylmethylamino-6,8-diiodo-2-(1-imidazolyl)quinazoline,
4-(2-phenoxyethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
6-hydroxymethyl-4-phenylmethylamino-2-(3-pyridyl)quinazoline,
6-methylthio-4-phenylmethylamino-2-(3-pyridyl)quinazoline,
6-methylsulfinyl-4-phenylmethylamino-2-(3-pyridyl)quinazoline,
6-methylsulfinyl-4-phenylmethylamino-2-(3-pyridyl)quinazoline,
4-phenylmethylamino-2-(2-thienyl)quinazoline,
4-phenylmethylamino-2-(2-furyl)quinazoline,
4-phenylmethylamino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline,
6-carboxy-4-phenylmethylamino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline,
6-ethoxycarbonyl-4-phenylmethylamino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline,
6-ethylaminocarbonyl-4-phenylmethylamino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline,
4-(2-methoxyethyl)amino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline or
4-(2-(2-hydroxyethoxy)ethyl)amino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline.

European published application number 0636626, which discloses compounds of the formula



and salts and solvates (e.g. hydrates) thereof, in which:

R¹ represents arylmethyl or C₁₋₆ alkyl optionally substituted by one or more fluorine atoms;

R² represents methyl;

R³ represents C₂₋₄ alkyl;

R⁴ represents nitro, cyano, C₁₋₆ alkoxy, C(=X)NR⁵R⁷, NR⁸R⁹, (CH₂)_mNR¹⁰C(=Y)R¹¹ or a 5-membered heterocyclic ring selected from thienyl, thiazolyl and 1,2,4-triazolyl each ring optionally substituted by a C₁₋₄ alkyl or aryl group; or when R¹ is arylmethyl or C₁₋₆ alkyl substituted by one or more fluorine atoms then R⁴ may also represent hydrogen;

R⁵ represents hydrogen or C₁₋₆ alkyl;

R⁶ represents hydrogen or C₁₋₆ alkyl;

R⁷ represents hydrogen, amino, hydroxyl, C₁₋₆ alkyl, aryl or arylC₁₋₄ alkyl;

R⁸ represents hydrogen or C₁₋₆ alkyl;

R⁹ represents hydrogen, C₁₋₆ alkyl, SO₂R¹², CO₂R¹², C(=NCN)SR¹² or C(=NCN)NR¹³R¹⁴;

R¹⁰ represents hydrogen or C₁₋₆ alkyl;

R¹¹ represents C₁₋₆ alkyl optionally substituted by one or more halogen atoms, or R¹¹ represents aryl, arylC₁₋₄ alkyl, thienyl, NR¹⁵R¹⁶, CH₂NR¹⁷R¹⁸ or R¹⁰ and R¹¹ together represent -A(CH₂)_n-;

R¹² represents C₁₋₆ alkyl, aryl or arylC₁₋₄ alkyl;

R¹³ represents hydrogen or C₁₋₆ alkyl;

R¹⁴ represents hydrogen, C₁₋₆ alkyl, aryl, arylC₁₋₄ alkyl or R¹³ and R¹⁴ together with the nitrogen atom to which they are attached form a morpholine, piperazine or N-C₁₋₄ alkylpiperazine ring;

R¹⁵ represents hydrogen or C₁₋₆ alkyl or R¹⁰ and R¹⁵ together represent -A(CH₂)_n-;

R¹⁶ represents hydrogen, C₁₋₆ alkyl, aryl, arylC₁₋₄ alkyl, CO₂R¹², CH₂CO₂R¹² or R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form a morpholine, piperazine or N-C₁₋₄ alkylpiperazine ring;

R¹⁷ represents hydrogen or C₁₋₆ alkyl;

R¹⁸ represents hydrogen, C₁₋₆ alkyl, aryl, arylC₁₋₄ alkyl, COR¹² or R¹⁷ and R¹⁸ together with the nitrogen atom to which they are attached form a morpholine, piperazine or N-C₁₋₄ alkylpiperazine ring;

A represents CH₂ or C=O;

m represents zero or 1;

n represents 1, 2 or 3;

X represents S or NH, or when R⁷ represents amino then X may also represent O;

Y represents O or S; for use in therapy.

Preferred compounds include:

1,3-Dimethyl-6-(2-propoxy-5-acetamidophenyl)-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-one;

1-ethyl-3-methyl-6-[2-propoxy-5-(4-methyl-2-thiazolyl)phenyl]-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-one;

1-ethyl-3-methyl-6-[2-propoxy-5-(2-methyl-4-thiazolyl)phenyl]-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-one;

1-ethyl-3-methyl-6-[2-propoxy-5-(2-(3-pyridyl)-4-thiazolyl)phenyl]-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-one;


1,3-dimethyl-6-[2-propoxy-5-(2-methyl-4-thiazolyl)phenyl]-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-one;

1,3-dimethyl-6-[2-propoxy-5-(3-phenyl-1,2,4-triazol-5-yl)phenyl]-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-one;

1,3-dimethyl-6-(2-propoxy-5-methanesulfonamidophenyl)-1,5-dihydro-pyrazolo[3,4-d]pyrimidin-4-one;

and physiologically acceptable salts and solvates (e.g. hydrates) thereof.

European published application number 0640599, which discloses compounds of the formula



wherein A is a bond, C1-4 alkylene or C1-4 oxyalkylene;

Y is a bond, C1-4 alkylene, C1-4 alkyleneoxy, C1-4 alkoxyphenylene or phenyl(C1-4)alkylene;

Z is a bond or vinylylene;

R1 is 4-15 membered heterocyclic ring containing one or two nitrogen atoms optionally substituted by one or two groups chosen from C1-4 alkyl, C1-4 alkoxy, halogen, trifluoromethyl and nitro;

R2 is (i) 4-15 membered heterocyclic ring containing one or two hetero atoms chosen from nitrogen, oxygen, and sulphur, not more than one hetero atom being sulphur, optionally substituted by one or two groups chosen from C1-4 alkyl, C1-4 alkoxy, halogen, trifluoromethyl, nitro and groups of formula:



wherein R10 is hydrogen or C1-4 alkyl,

(ii) C4-15 carbocyclic ring,

(iii) C1-4 alkoxy,

(iv) hydroxy(C1-4 alkoxy) or

(v) hydroxy;


R3 is (i) 4-15 membered heterocyclic ring containing one or two hetero atoms chosen from nitrogen, oxygen and sulphur, not more than one hetero atom being oxygen or sulphur, optionally substituted by one or two groups chosen from C1-4 alkyl, C1-4 alkoxy, halogen, trifluoromethyl, nitro, cyano, ethynyl and groups of formula:



wherein R7 and R8 are independently hydrogen or C1-4 alkyl.

(ii) C4-15 carbocyclic ring,

(iii) a group of formula:




wherein X is halogen, or

(iv) hydrogen,

and I is 1 or 2,

provided that: R2 is not hydroxy when Y is a bond; R1 is not bonded through its nitrogen atom when Z is vinylylene; and excluding compounds of the formula:



wherein RAA is methyl or n-propyl;


RBB is cyclopentyl, cyclohexyl, 2-hydroxyethyl, methoxyethyl, 2-(1-piperidinyl)ethyl, or phenyl or benzyl which may be substituted by 1 or 2 of methyl, methoxy, chloro, nitro and trifluoromethyl;

RCC is hydrogen or methyl;

RDD is methyl or n-propyl, isopropyl or benzyl; and

REE is hydrogen or methyl;

and the compound of formula:




and its pharmaceutically acceptable salts.

Preferred compounds include:

2-(1-Imidazolyl)-4-[2-(2-hydroxyethoxy)ethyl]amino-5-(3-methoxyphenyl)-methylpyrimidine,
2-(1-Imidazolyl)-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-4-(2-methoxyethyl)aminopyrimidine,
2-(1-Imidazolyl)-5-ethyl-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-5-phenylmethyl-4-phenylmethylaminopyrimidine
2-(1-Imidazolyl)-5-methyl-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-5,6-dimethyl-4-phenylmethylaminopyrimidine


2-(1-Imidazolyl)-5-(3-methoxyphenyl)methyl-4-(2-methoxyethyl)amino-pyrimidine,
2-(1-Imidazolyl)-5-(4-methoxyphenyl)methyl-4-[2-(2-hydroxyethoxy)ethyl]aminopyrimidine,
2-(1-Imidazolyl)-5-(4-methoxyphenyl)methyl-4-(2-methoxyethyl)amino-pyrimidine,
2-(1-Imidazolyl)-5-(4-methoxyphenyl)methyl-4-phenylmethylamino-pyrimidine,
2-(1-Imidazolyl)-5-phenoxymethyl-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-5-(1-imidazolyl)methyl-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thiazolyl)-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(1,3-dioxalindan-5-yl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-[2-(2-hydroxyethoxy)ethyl]aminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(1-naphthyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(4-methoxyphenyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(3-methoxyphenyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(2-furyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(2-thienyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(3-pyridyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(2-methoxyethyl)aminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-phenylmethoxyaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(4-chlorophenyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(3-chlorophenyl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(1,3-dioxalindan-5-yl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(4-methylphenyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine,
2-(1-Imidazolyl)-5-(4-methoxyphenyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine,
2-(1-Imidazolyl)-5-(5-methyl-2-thienyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-[4-(1-imidazolyl)phenyl]methylamino-pyrimidine,
2-(1-Imidazolyl)-5-(3-pyridyl)-4-(1,3-dioxalindan-5-yl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(3-furyl)-4-(1,3-dioxalindan-5-yl)methylaminopyrimidine,
2-(1-Imidazolyl)-5-(3-pyridyl)-4-phenylmethylaminopyrimidine,
2-(1-Imidazolyl)-5-(4-chlorophenyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine,
2-(Benzimidazol-1-yl)-5-(2-thienyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine,
2-(1-Imidazolyl)-5-(2-thienyl)-4-(4-ethoxycarbonylphenyl)methylamino-pyrimidine,
2-(1-Imidazolyl)-5-(2-naphthyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine,
2-(3-Pyridyl)-5-(2-thienyl)-4-(1,3-dioxalindan-5-yl)methylaminopyrimidine,
2-[2-(3-Pyridyl)vinyl]-5-(2-thienyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine,
2-(2-Methyl-1-imidazolyl)-5-(2-thienyl)-4-(1,3-dioxalindan-5-yl)methylamino-pyrimidine or
2-(1-Imidazolyl)-5-(2-thienyl)-4-(benzimidazol-5-yl)methylaminopyrimidine

European published application number 0668280, which discloses compounds of the formula



wherein R¹ and R² are the same or different and represent hydrogen, lower alkyl (which is optionally substituted with one to three substituents which are the same or different and are cycloalkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, halogen, alicyclic heterocycle group (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, aralkyl, aryl optionally substituted with one to three substituents which are the same or different and are lower alkoxy, or aromatic heterocycle group)), cycloalkyl, bicycloalkyl, benzocycloalkyl (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, sulfonamide, halogen, or trifluoromethyl), lower alkenyl, aryl (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, sulfonamide, halogen, or trifluoromethyl), aromatic heterocycle group-substituted alkyl (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, sulfonamide, halogen or trifluoromethyl and where said alkyl part is optionally substituted with aryl), aromatic heterocycle group (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, sulfonamide, halogen, or trifluoromethyl), or aralkyl (where the aryl part of said aralkyl is optionally substituted with one to three substituents which are the same or different and are lower alkyl, lower alkoxy, dialkyl-substituted amino, halogen, or trifluoromethyl), or R¹ and R² are taken together to represent heterocycle group containing nitrogen atom (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, aryl, or aralkyl), R³ represents hydrogen, lower alkyl (which is optionally substituted with one to three substituents which are the same or different and are cycloalkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, halogen, or alicyclic heterocycle group (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, aralkyl, aryl optionally substituted with one to three substituents which are the same or different and are lower alkoxy, or aromatic heterocycle group)), cycloalkyl, lower alkenyl, aryl (which is optionally substituted with one to three substituents which are the same or different and are lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, sulfonamide, halogen, or trifluoromethyl), aromatic heterocycle group-substituted alkyl (where said aromatic heterocycle group part is optionally substituted with one to three substituents which are the same or different and are lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, sulfonamide, halogen or trifluoromethyl, and where the alkyl part is optionally substituted with aryl), aromatic heterocycle group (where said aromatic heterocycle group is optionally substituted with one to three substituents which are the same or different and are lower alkyl, hydroxy, lower alkoxy, carboxy, lower alkoxy carbonyl, amino, monoalkyl-substituted amino, dialkyl-substituted amino, nitro, sulfonamide, halogen, or trifluoromethyl), or aralkyl (where the aryl part of said aralkyl is optionally substituted with one to three substituents which are the same or different and are lower alkyl, lower alkoxy, dialkyl-substituted amino, halogen, or trifluoromethyl), and X represents oxygen atom or sulfur atom, or pharmacologically acceptable salts thereof.

European published application number 0669324, which discloses compounds of the formula




(wherein R^1 , R^2 , R^3 , R^4 and R^5 may be the same or different from each other and each represents a hydrogen atom, a halogen atom, a lower alkyl group or a lower alkoxy group; and

R^6 and R^7 may be the same or different from each other and each represents a hydrogen atom, a lower alkyl group, a hydroxyalkyl group, a lower alkoxyalkyl group, a cyanoalkyl group, a heteroarylalkyl group, a cycloalkyl group, a cycloalkylalkyl group or a carboxyl alkyl group which may be protected, or alternatively R^6 and R^7 may form a ring together with the nitrogen atom to which they are bonded, this ring optionally having a substituent).

or a pharmacologically acceptable salt thereof:

WO91/19717 discloses compounds of the formula.



wherein

J is oxygen or sulfur,

R^1 is hydrogen, alkyl or alkyl substituted with aryl or hydroxy;

R^2 is hydrogen, aryl, heteroaryl, cycloalkyl, alkyl or alkyl substituted with aryl, heteroaryl, hydroxy, alkoxy, amino, monoalkyl amino or dialkylamino, or $-(CH_2)_mTCOR^{20}$ wherein m is an integer from 1 to 6, T is oxygen or $-NH-$ and R^{20} is hydrogen, aryl, heteroaryl, alkyl or alkyl substituted with aryl or heteroaryl;

R³ is hydrogen, halo, trifluoromethyl, alkoxy, alkylthio, alkyl, cycloalkyl, aryl, aminosulfonyl, amino, monoalkylamino, dialkylamino, hydroxyalkylamino, aminoalkylamino, carboxy, alkoxycarbonyl or aminocarbonyl or alkyl substituted with aryl, hydroxy, alkoxy, amino, monoalkylamino or dialkylamino;

R^a, R^b, R^c and R^d independently represent hydrogen, alkyl, cycloalkyl or aryl; or (R^a and R^b) or (R^c and R^d) or (R^b and R^c) can complete a saturated ring of 5- to 7- carbon atoms, or (R^a and R^b) taken together and (R^b and R^c) taken together, each complete a saturated ring of 5- to 7-carbon atoms, wherein each ring optionally can contain a sulfur or oxygen atom and whose carbon atoms may be optionally substituted with one or more of the following: alkenyl, alkynyl, hydroxy, carboxy, alkoxycarbonyl, alkyl or alkyl substituted with hydroxy, carboxy or alkoxycarbonyl; or such saturated ring can have two adjacent carbon atoms which are shared with an adjoining aryl ring; and
n is zero or one.

Preferred compounds include:

cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(phenylmethyl)-
cyclopenta[4,5]imidazo[2,1-b]purin-4-one;
7,8-Dihydro-5-methyl-3-(phenylmethyl)-3*H*-imidazo[2,1-b]purin-4(5*H*)-
one;
cis-6a,7,8,9,10,10a-Hexahydro-5-methyl-3-(phenylmethyl)-3*H*-
benzimidazo[2,1-b] purin-4(5*H*)-one;
5,7,8,9-Tetrahydro-5-methyl-3-(phenylmethyl)pyrimido[2,1-b]purin-
4(3*H*)-one;
7,8-Dihydro-8-phenyl-5-methyl-3-(phenylmethyl)-3*H*-imidazo[2,1-
b]purin-4(5*H*)-one;
5',7'-Dihydro-5'-methyl-3'-(phenylmethyl)spiro[cyclohexane-1,8'-(8*H*)-
imidazo[2,1-b]purin]-4'(3'*H*)-one;
cis-5,6a,11,11a-Tetrahydro-5-methyl-3-
(phenylmethyl)indeno[1',2':4,5]imidazo[2,1-b]purin-4(3*H*)-one;
5',7'-Dihydro-2',5' dimethyl-3'-(phenylmethyl)spiro{cyclohexane-
1,7'(8'*H*)-imidazo[2,1-b]purin}-4'(3'*H*)-one;
7,8-Dihydro-2,5,7,7,8(R,S)-pentamethyl-3*H*-imidazo[2,1-b]purin-4(5*H*)-
one;
cis-5,6a,7,11b-Tetrahydro-5-methyl-3-

(phenylmethyl)indeno[2',1':4,5]imidazo[2,1-b]purin-4(3*H*)-one;
cis-5,6a,7,8,9,9a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-
cyclopent[4,5]imidazo[2,1-b]purin-4-(3*H*)-one;
5'-Methyl-3'-(phenylmethyl)-spiro[cyclopentane-1,7'(8'*H*)-(3'*H*)-
imidazo[2,1-b]purin]-4'(5'*H*)-one;
7,8-Dihydro-2,5,7,7-tetramethyl-3-(phenylmethyl)-3*H*-imidazo[2,1-
b]purin-4(5*H*)-one;
7,8-Dihydro-7(*R*)-phenyl-2,5-dimethyl-3-(phenylmethyl)-3*H*-imidazo[2,1-
b]purin-4(5*H*)-one;
7,8-Dihydro-2,5-dimethyl-3,7(*R*)-bis(phenylmethyl)-3*H*-imidazo[2,1-
b]purin-4(5*H*)-one;
(±)-7,8-Dihydro-2,5-dimethyl-7-ethyl-3-(phenylmethyl)-3*H*-imidazo[2,1-
b]purin-4(5*H*)-one;
6a(*S*)-7,8,9,10,10a(*R*)-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-3*H*-
benzimidazo[2,1-b]purin-4(5*H*)-one;
6a(*R*)-7,8,9,10,10a(*S*)-hexahydro-2,5-dimethyl-3-(phenylmethyl)-3*H*-
benzimidazo[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5-dimethyl-7(*R*)-isopropyl-3-(phenylmethyl)-3*H*-
imidazo[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5,7(*R*)-trimethyl-3-(phenylmethyl)-3*H*-imidazo[2,1-b]purin-
4(5*H*)-one;
cis-7,7a,8,9,10,10a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-3*H*-
cyclopenta[5,6]pyrimido[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5-dimethyl-7(*S*)-(1-methylpropyl)-3-(phenylmethyl)-3*H*-
imidazo[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5-dimethyl-7(*R*)-(2-methylpropyl)-3-(phenylmethyl)-3*H*-
imidazo[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5-dimethyl-7(*R,S*)-(methoxycarbonyl)-3-(phenylmethyl)-
3*H*-imidazo[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5-dimethyl-7(*R,S*)-(1-propyl)-3-(phenylmethyl)-3*H*-
imidazo[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5-dimethyl-7(*S*)-(1-methylethyl)-3-(phenylmethyl)-3*H*-
imidazo[2,1-b]purin-4(5*H*)-one;
7,8-Dihydro-2,5,7,7,8(*R,S*)-pentamethyl-3*H*-imidazo[2,1-b]purin-4(5*H*)-
one;
5,7,8,9-Tetrahydro-2,5,7,9(*R,S*)-pentamethyl-3-(phenylmethyl)-
pyrimido[2,1-b]purin-4(3*H*)-one;
5,6a(*R*),7,8,9,9a(*S*)-Hexahydro-2,5-dimethyl-3-
(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
5,6a(*S*),7,8,9,9a(*R*)-Hexahydro-2,5-dimethyl-3-
(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;


cis-6a,7,8,9,10,10a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
5',7'-Dihydro-2',5'-dimethyl-3'-(phenylmethyl)spiro[cyclohexane-1,8'-(8H)-imidazo[2,1-b]purin]-4'(3'H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-cyclohept[6,7]imidazo[2,1-b]purin-4(3H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-ethyl-3-(phenylmethyl)-cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
cis-6a,7,8,9,10,10a-Hexahydro-5-methyl-2-ethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-ethyl-3-(phenylmethyl)-cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-phenyl-3-(phenylmethyl)-cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
cis-6a,7,8,9,10,10a-Hexahydro-5-methyl-2-phenyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-5-methylcyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-2,5-dimethylcyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-one;
cis-5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-di-methyl-cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
2'-Methyl-3'-spiro[cyclopentane-1,7'(8'H)-(3'H)-imidazo[2,1-b]purin]-4'(5'H)-one;
7,8-Dihydro-2,5-dimethyl-7(R)-(1-methylethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
7,8-Dihydro-2,5,7,7-tetramethyl-3H-imidazo[2,1-b]purin-4(5H)-one;
7,8-Dihydro-2,5-dimethyl-7(S)-(1-methylethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
6a(R),7,8,9,10,10a(S)-Hexahydro-2,5-dimethyl-3H-benzimidazo[2,1-b]purin-4(5H)-one;
5',7'-Dihydro-2',5'-dimethylspiro[cyclohexane-1,7'(8'H)-imidazo[2,1-b]purin]-4'(3'H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(phenylmethyl)cyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-thione;
5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-thione;
cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(4-chlorophenylmethyl)cyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-one;
cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(cyclohexylmethyl)-

cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(2-naphthylmethyl)-
 cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 bromophenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 5,6a(R)-7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(4-
 methoxyphenylmethyl)-cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-
 one;
 cis-5,6a,7,8,9,9a-Hexahydro-2,3,5-trimethylcyclopent[4,5]imidazo[2,1-
 b]purin-4(3*H*)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2-(hydroxymethyl)-5-methyl-3-
 (phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2-methylthio-5-methyl-3-(phenylmethyl)-
 cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 cis-3,4,5,6a,7,8,9,9a-Octahydro-5-methyl-4-oxo-3-(phenylmethyl)-
 cyclopent[4,5]imidazo[2,1-b]purin-2-carboxylic acid;
 cis-3,4,5,6a,7,8,9,9a-Octahydro-5-methyl-4-oxo-3-(phenylmethyl)-
 cyclopent[4,5]imidazo[2,1-b]purin-2-carboxylic acid, methyl ester;
 cis-5,6a,7,8,9,9a-Hexahydro-2-bromo-5-methyl-3-(phenylmethyl)-
 cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2-(methylaminosulfonyl)-5-methyl-3-
 (phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 cis-1-Cyclopentyl-5,6a,7,8,9,9a-hexahydro-5-methyl-
 cyclopent[4,5]imidazo[2,1-b]purin-4(1*H*)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-3,5-bis-(phenylmethyl)
 cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 cis-6a,7,8,9,10,10a-Hexahydro-3,5-bis-(phenylmethyl)-3*H*-
 benzimidazo[2,1-b]purin-4(5*H*)-one;
 cis-3-Cyclopentyl-5,6a,7,8,9,9a-hexahydro-5-methyl-
 cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 5'-Methyl-3'-(phenylmethyl)spiro[cyclopentane-1,7'-(8'*H*)-(3'*H*)-
 imidazo[2,1-b]purin]-4'(5'*H*)-one;
 2',5'-Dimethyl-3'-(phenylmethyl)-spiro[cyclopentane-1,7'-(8'*H*)-(3'*H*)-
 imidazo[2,1-b]purin]-4'(5'*H*)-one;
 cis-5,6a,(R)7,8,9,9a(S)-Hexahydro-5-methyl-3-
 (phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;
 cis-3-Cyclopentyl-5,6a,7,8,9,9a-Hexahydro-2,5-
 dimethylcyclopent[4,5]imidazo[2,1-b]purin-4(3*H*)-one;36
 5'-Methyl-2'-trifluoromethyl-3'-(phenylmethyl)spiro[cyclopentane-
 1,7'-(8'*H*)-(3'*H*)-imidazo[2,1-b]purin]-4'(5'*H*)-one;
 7,8-Dihydro-5,7,7-trimethyl-2-trifluoromethyl-3-(phenylmethyl)-3*H*-
 imidazo[2,1-b]purin-4(5*H*)-one;

- (+/-)-cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-trifluoromethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
- (+/-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3-(phenylmethyl)-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)-one;
- (+)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3-phenylmethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)-one;
- (-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3-phenylmethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)-one;
- (+/-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)-one;
- (+)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)-one;
- (-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)-one;
- 6a,7,8,9,10,10a,11,12,13,13a-Decahydro-2,5-dimethyl-(3-phenylmethyl)naph[1,8a-d]imidazo[2,1-b]purin-4(5H)-one;
- 7(R)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(3H)-one;
- 7(R)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3H-imidazo[2,1-b]purin-4(5H)-one;
- 7(S)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(3H)-one;
- 7(S)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3H-imidazo[2,1-b]purin-4(5H)-one;
- 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-[(trimethylacetoxymethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one];
- 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(4-pyridylmethyl)-cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
- 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-[2-(1-morpholinyl)ethyl]cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
- 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-[acetoxymethyl]cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
- 5,6a,7,8,9,9a-Hexahydro-2,5,6a-trimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
- 5,6a(R),7(S),8,9,9a-Hexahydro-2,5,6a-trimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
- 5,6a(S),7(R),8,9,9a-Hexahydro-2,5,6a-trimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
- cis-6a,7,8,9,10,10a-Hexahydro-2,5,7-trimethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;

cis-5,6a,7,8,9,9a-Hexahydro-2,5,6a-trimethylcyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one]; or
cis-6a,7,8,9,10,10a-Hexahydro-2,5,7-trimethyl-3H-benzimidazo[2,1-b]purin-4(5H)-one].

WO 94/19351 discloses compounds of the formula



or a pharmaceutically acceptable salt thereof, wherein:

R₁, R₂ and R₃ are independently selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, halogeno, hydroxy, (di-lower alkyl)amino, 4-morpholinyl, 1-pyrrolidinyl, 1-pyrrolyl, -CF₃, -OCF₃, phenyl and methoxyphenyl; or R₁ and R₂ together are methylenedioxy; or R₁ and R₂ together with the carbon atoms to which they are attached form a benzene ring; and

R^a is hydrogen and R^b and R^c, together with the carbon atoms to which they are attached, form a saturated ring of 5 carbons; or R^a is lower alkyl, R^b is hydrogen or lower alkyl, and R^c is hydrogen; or R^a, R^b and the carbon atom to which they are attached form a saturated ring of 5-7 carbons, and R^c is hydrogen; or R^a is hydrogen, and R^b, R^c and the carbon atoms to which they are attached form a tetrahydrofuran ring; or R^a and R^b, together with the carbon atom to which they are attached, and R^b and R^c, together with the carbon atoms to which they are attached, each form a saturated ring of 5-7 carbons.

-39-

Preferred compounds include:

2'-benzyl-spiro[cyclopentane-1',7' (8'H)-[3'H]-imidazo[2,1-b]purin-4'-(5'H)-one;

2'-benzyl-5,7,7-trimethyl-3H-imidazo[2,1-b]purin-4-(5H)-one;


(+)-2-benzyl-7, 8-dihydro-5-methyl-7-(1-methylethyl)-1H-imidazo[2,1-b]-purin-4(5H)-one;

(+,-)-6a, 7, 8, 9, 9a, 10, 11, 11a-octahydro-5-methyl-2-(3,4-methylene-dioxyphenylmethyl)-3H-pentalen[6a,1:4,5]imidazo[2,1-b]purin-4(5H)-one; and

(+)-cis-6a, 7, 9, 9a-tetrahydro-5-methyl-2-[4-(trifluoromethyl)-phenylmethyl]-3H-furo[3', 4':4,5]imidazo[2,1-b]purin-4(5H)-one.

WO 94/22855 discloses compounds of the formula

1. A nitrogen-containing fused-heterocyclic compound having the formula (I) or a pharmacologically acceptable salt thereof:




in which ring A represents a benzene, pyridine or cyclohexane ring and B represents a pyridine, imidazole or pyrimidine ring, with the proviso that rings A and B are bonded to each other with two atoms being shared by them, and the shared atoms may be any of carbon and nitrogen atoms;

R^1 represents a group represented by the formula: $-NR^4R^5$ (wherein R^4 and R^5 may be the same or different

-40-


from each other and each represent a hydrogen atom, a lower alkyl or acyl group or a carboxyl group which may be protected, or alternatively R^4 and R^5 may form a ring together with the nitrogen atom to which they are bonded, provided that the ring may be substituted), or a heteroaryl group which has one or two nitrogen atoms and may be substituted;

R^2 represents a hydrogen atom, a group represented by the formula:



(wherein R^8 represents a carboxyl or tetrazolyl group which may be protected),
or a halogen atom;
and


R^3 represents a hydrogen atom or a group represented by the formula:



(wherein R^6 and R^7 each represent a hydrogen or halogen atom or a lower alkoxy group, or alternatively R^6 and R^7 may together form a methylenedioxy or ethylenedioxy group).

-41-

WO 95/19978 discloses compounds of the formula




and salts and solvates thereof, in which:

R^0 represents hydrogen, halogen or C_{1-6} alkyl;

R^1 represents hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl C_{1-3} alkyl, aryl C_{1-3} alkyl or heteroaryl C_{1-3} alkyl;

R^2 represents an optionally substituted monocyclic aromatic ring selected from benzene, thiophene, furan and pyridine or an optionally




substituted bicyclic ring attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen; and R^3 represents hydrogen or C_{1-3} alkyl, or R^1 and R^3 together represent a 3- or 4- membered alkyl or alkenyl chain.

Preferred compounds include:

Cis-2,3,6,7,12,12a-hexahydro-2-(4-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2', 1' : 6, 1]pyrido[3,4-b]indole-1,4-dione;
 Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methyl-pyrazino[2', 1' : 6, 1]pyrido[3,4-b]indole -1,4-dione;
 Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-methyl-pyrazino[2', 1' : 6, 1]pyrido[3,4-b]indole -1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (5aR,12R,14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1",2" : 4',5']pyrazino[2',1' : 6,1]pyrido[3,4-b]indole-5-1,4-dione;
 and physiologically acceptable salts and solvates thereof.

U.S. Patent No. 5,294,612 discloses compounds of the formula



wherein:

R¹ is hydrogen, alkyl, C₄ to C₇ cycloalkyl, C₄ to C₇ cycloalkyl substituted by C₁ to C₁₀ alkyl or hydroxyl, 2- or 3-tetrahydrofuranyl, 3-tetrahydrothienyl 1,1, -dioxide, C₄ to C₇ cycloalkyl-C₁ to C₁₀ alkyl, carboxy-C₁ to C₁₀ alkyl, carbo-C₁ to C₄ lower-alkoxy-C₁ to C₁₀ alkyl, dialkylamino C₁ to C₁₀ alkyl, phenyl-C₁ to C₄ lower-alkyl, phenyl-C₁ to C₄ lower-alkyl in which the phenyl ring is substituted in the 2, 3, or 4-position by one or two substituents, the same or different, selected from the group consisting of amino, halogen, C₁ to C₁₀ alkyl, carboxyl, carbo-C₁ to C₄ lower-alkoxy, carbamoyl, NHSO₂ (quinolinyl), nitro and cyano;


R³ is, C₁ to C₄ lower-alkyl, phenyl-C₁ to C₄ lower-alkyl, lower-alkoxyphenyl-C₁ to C₄ lower-alkyl, diC₁ to C₄ lower-alkoxy-phenyl-C₁ to C₄ lower-alkyl, pyridyl-C₁ to C₄ lower-alkyl, C₄ to C₇ cycloalkyl-C₁ to C₄ lower-alkyl, phenylamino, diC₁ to C₁₀ alkylamino, halogen, trifluoromethyl, C₁ to C₄ lower-alkylthio, cyano or nitro; and

R⁶ is a nine or ten membered bicyclic ring having carbon and from one to two nitrogen atoms, and

the heterocycle is made up of fused 5 or 6 membered rings or such ring substituted at any available carbon atom by one or two substituents, the same or different, selected from the group consisting of C₁ to C₄ lower-alkyl, halogen, C₁ to C₄ lower-alkoxy, C₄ to C₇ cycloalkyloxy, 4-morpholinyl, C₁ to C₄ lower-alkoxy-C₁ to C₄ lower-alkoxy, hydroxy, imidazolyl, oxo and 4-morpholinyl-C₁ to C₄ lower-alkoxy, or at any available nitrogen atom by C₁ to C₄ lower-alkyl, C₂ to C₄ lower-alkanoyl, or trifluoroacetyl; or a pharmaceutically acceptable acid-addition salt thereof.

formula

U.S. Patent No. 5,405,847 discloses compounds of the



where the benzo ring can also contain a nitrogen atom instead of a CH group either in position 6, 7, 8 or 9 and the radicals R₁, R₂, R₃ and R₄ have the following meanings:

R₁: C₂-C₆-alkenyl, C₂-C₆-alkynyl, hydroxy, C₁-C₆-alkoxy, C₃-C₆-alkenylloxy, C₃-C₆-alkynylloxy, C₂-C₆-alkanoyloxy, benzoyloxy, morpholinocarbonyloxy, C₁-C₆-alkyloxy carbonyloxy, C₁-C₆-alkylaminocarbonyloxy, C₁-C₆-dialkylaminocarbonyloxy or the group

-Akk-A

where Alk: is C₁-C₆-alkyl, C₂-C₆-hydroxyalkyl or C₃-C₆-cycloalkyl and the symbol A represents:

- 1) Hydrogen, halogen, hydroxy, C₁-C₆-alkoxy, C₂-C₆-alkanoyloxy, phenyl;
- 2) -NHR₅, -NR₅R₆, NR₅R₆R₇, pyridylamino, imidazolyl, pyrrolidinyl, N-C₁-C₆-alkylpyrrolidi-

nyl, piperidylamino, N-(phenyl-C₁-C₆-alkyl)-piperidylamino where R₅ and R₆ may be the same or different and represent hydrogen, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, C₃-C₇-hydroxycycloalkyl, morpholino-C₁-C₆-alkyl, phenyl, phenyl-C₁-C₆-alkyl or phenyl-C₂-C₆-oxyalkyl, it also being possible for the phenyl radicals in R₅ and R₆ to be substituted by halogen and R₇ is hydrogen or C₁-C₆-alkyl;


- 3) The group:

CD.D

where D is phenyl, C₁-C₆-alkyl, C₃-C₇-cycloalkyl, hydroxy, C₁-C₆-alkoxy, C₃-C₇-cycloalkyloxy, morpholino, pyrrolidino, piperidino, homopiperidino, piperazino, -NHR₅ or -NR₅R₆ and R₅ and R₆ have the meanings given herein-above.


-44-

4) The group:




where n can be the integers 1-3 and E represents CH₂, oxygen, sulfur, NH, CHOH, CH—C₁—C₆—alkyloxy, CH—C₂—C₆—alkanoyloxy, CHC₆H₅, CHCO₂D, CH—CH₂C₆H₅, N—C₁—C₆—alkyl, N—C₁—C₆—hydroxyalkyl, N—C₆H₅, N—CH₂C₆H₅, N—CH(C₆H₅)₂, N—(CH₂)₂—OH, N—(CH₂)₃—OH or NCOD and the phenyl radicals (C₆H₅) may also be substituted by halogen, C₁—C₆—alkoxy, trifluoromethyl, C₁—C₆—alkyl, methylenedioxy or cyan and D has the meanings given hereinabove;

R₂ and R₃, which may be the same or different: hydrogen, halogen, hydroxy, C₁—C₆—alkyl, trifluoromethyl, —CN, C₁—C₆—alkoxy, C₃—C₆—alkenyloxy, C₃—C₆—alkynyloxy, —NHR₅, —NR₅R₆, —NR₅R₆R₇ (meanings R₅, R₆, R₇ as given hereinabove) or the group —G-Alk-A, where Alk and A have the meanings given hereinabove and G is oxygen, sulfur, NH or NR₅ and R₂ can also be




R₄: hydrogen or halogen, where R₁ can also be hydrogen, when R₂ is the group



and R₅ represents phenyl, C₁—C₄—alkoxyphenyl or diphenylmethyl and R₁ and R₄ are hydrogen, and their physiologically acceptable acid addition salts and quaternary ammonium salts, with the exception of the compounds of Formula I where R₁ is methyl, dimethylaminopropyl, dimethylaminoethyl, morpholinoethyl or pyrrolidinoethyl, R₂, R₃ and R₄ are hydrogen and the benzo ring does not contain a nitrogen atom instead of a CH group.

U.S. Patent No. 5,436,233 discloses compounds of the

formula



wherein R¹ is hydrogen or C1-4 alkyl;

Y is single bond or C1-6 alkylene;

A is

(i) —CyA—(R²)_p,

(ii) —O—R⁰ or —S(O)_p—R⁰,

in which R⁰ is R^{0A} or R^{0B};

R^{0A} is —CyA—(R²)_p;

R^{0B} is hydrogen or C1-4 alkyl;

p is 0-2;

CyA is

(1) 3-7 membered, saturated or unsaturated, monocyclic carbocyclic ring,

- (2) 7-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one nitrogen atom, one nitrogen and one oxygen atoms, two nitrogen and one oxygen atoms, or one nitrogen and two oxygen atoms,
- (3) 6-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one nitrogen and one oxygen atoms, two nitrogen and one oxygen atoms, or one nitrogen and two oxygen atoms,
- (4) 6-membered, unsaturated or partially saturated, monocyclic hetero ring containing as a hetero atom, one nitrogen atom,
- (5) 4- or 5-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one nitrogen atom, one nitrogen and one oxygen atoms, two nitrogen and one oxygen atoms, or one nitrogen and two oxygen atoms,
- (6) 4-7 membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one or two sulfur atoms or
- (7) 4-7 membered, unsaturated or partially or fully saturated, monocyclic hetero ring containing as hetero atoms, one or two oxygen atom;

R^2 is R^{2A} or R^{2B} ;

R^{2A} is (1) $-NR^6AR^{7A}$, in which R^{6A} and R^{7A} independently are hydrogen or C1-4 alkyl (with the proviso that R^{6A} and R^{7A} are not hydrogen at same time), (2) $-SO_2NR^6R^7$, in which R^6 and R^7 independently are hydrogen or C1-4 alkyl, (3) trifluoromethyl or (4) trifluoromethoxy;

R^{2B} is (1) hydrogen, (2) C1-4 alkyl, (3) C1-4 alkoxy, (4) $-COOR^5$, in which R^5 is hydrogen or C1-4 alkyl, (5) halogen, (6) nitro or (7) $-NR^6BR^{7B}$, in which R^{6B} and R^{7B} are hydrogen;

Z is Z^A or Z^B ;

Z^A is methylene, ethylene, vinylene or ethynylene;

Z^B is single bond;

CyB is

- (1) 7-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one, two or three nitrogen atoms,
- (2) 6-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, two or three nitrogen atoms,
- (3) 6-membered, unsaturated or partially saturated, monocyclic hetero ring containing as a hetero atom, one nitrogen atom,
- (4) 4- or 5-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one, two or three nitrogen atoms, or
- (5) 4-7 membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one or two oxygen atoms, or one or two sulfur atoms;

R^3 is hydrogen, C1-4 alkyl, C1-4 alkoxy, halogen or trifluoromethyl;

R^4 is R^{4A} or R^{4B} ;

R^{4A} is (1) $-NH-SO_2R^{11}$, in which R^{11} is C1-4 alkyl, (2) $SO_2NR^9R^{10}$, in which

R^9 is hydrogen, C1-4 alkyl or phenyl(C1-4 alkyl) and R^{10} is hydrogen or C1-4 alkyl, (3) $-OCOR^{11}$, in which R^{11} is as hereinbefore defined, (4) hydroxy, (5) $-SO_2N=CHNR^{12}R^{13}$ in which R^{12} is hydrogen or C1-4 alkyl and R^{13} is C1-4 alkyl, (6) $-CONR^{14}R^{15}$ in which R^{14} is hydrogen or C1-4 alkyl and R^{15} is C1-4 alkyl or phenyl(C1-4 alkyl), (7) ethynyl, (8) tri(C1-4 alkyl)silylethynyl or (9) acetyl;

R^{4B} is (1) hydrogen, (2) C1-4 alkyl, (3) C1-4 alkoxy, (4) $-\text{COOR}^8$, in which R^8 is hydrogen or C1-4 alkyl, (5) $-\text{NR}^9\text{R}^{10}$, in which R^9 and R^{10} are as hereinbefore defined, (6) $-\text{NHCOR}^{11}$, in which R^{11} is as hereinbefore defined, (7) halogen, (8) trifluoromethyl, (9) nitro, (10) cyano, (11) C1-4 alkylthio, (12) C1-4 alkylsulfinyl, (13) C1-4 alkylsulfonyl, (14) hydroxymethyl, and l, m and n independently are 1 or 2; with the proviso that

- (1) the group of the formula: $-\text{CyA}-(\text{R}^2)_l$ does not represent a cyclopentyl and trifluoromethylphenyl group when Y is a single bond, that
- (2) a CyB ring does not bond to Z through a nitrogen atom in the CyB ring when Z is vinylene or ethynylene, that
- (3) a CyB ring is not pyridine or thiophene when CyA is a ring of $\text{CyA}-(7)$ that
- (4) Y is not a single bond, when A is (ii) $-\text{O}-\text{R}^0$ or $-\text{S}(\text{O})_p-\text{R}^0$ and that
- (5) A is not $-\text{CyA}-(\text{R}^2\text{B})_l$ and $-\text{OR}^{0B}$, when Z is Z^B and R^4 is R^{4B} ; or pharmaceutically acceptable acid addition salts thereof, pharmaceutically acceptable salts thereof, or hydrates thereof.

Preferred compounds include:

4-phenylmethylamino-2-((1-imidazolyl)methyl)-quinazoline,
 4-phenylmethylamino-2-((1-imidazolyl)methyl)-quinazoline,
 6-chloro-4-phenylmethylamino-2-(1-imidazolylmethyl)-quinazoline,
 6-chloro-4-phenylamino-2-(1-imidazolylmethyl)-quinazoline,
 6-chloro-4-(3-carboxyphenyl)amino-2-(1-imidazolylmethyl)-quinazoline
 or
 4-phenylmethylamino-2-(2-(3-pyridyl)vinyl)quinazoline,
 and pharmaceutically acceptable acid addition salts thereof, pharmaceutically acceptable salts thereof, or hydrates thereof.


6-dimethylaminosulfonyl-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
 6-dimethylaminomethylideneaminosulfonyl-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
 6-(phenylmethylaminosulfonyl)-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
 6-phenylmethylaminocarbonyl-4-phenylmethylamino-2-(1-imidazolyl)quinazoline,
 6-ethylaminocarbonyl-4-phenylmethylamino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline,
 6-hydroxy-4-phenylmethylamino-2-(1-imidazolyl)-quinazoline,
 6-(1-imidazolyl)-4-(2-methoxyethyl)amino-6-(2-triethylsilylethynyl)quinazoline,
 6-ethynyl-4-(2-methoxyethyl)amino-2-(1-imidazolyl)-quinazoline,
 6-(1-imidazolyl)-4-phenylmethylamino-6-ethynyl-quinazoline or
 6-acetyl-4-(2-methoxyethyl)amino-2-(1-imidazolyl)-quinazoline,
 and pharmaceutically acceptable acid addition salts thereof, pharmaceutically acceptable salts thereof, or hydrates thereof.

-47-


4-(2-methylthioethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
 4-(2-methylsulfinylethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
 4-(2-methylsulfonylethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline,
 4-(3-trifluoromethylphenylmethyl)amino-2-(3-pyridyl)quinazoline,
 4-(4-(N,N-dimethylamino)phenylmethyl)amino-2-(3-pyridyl)quinazoline,
 4-(4-sulfamoylphenylmethyl)amino-2-(3-pyridyl)quinazoline,
 4-(4-trifluoromethoxyphenylmethyl)amino-2-(1-imidazolyl)quinazoline,
 4-(3-trifluoromethoxyphenylmethyl)amino-2-(1-imidazolyl)quinazoline,
 4-(2-phenoxyethyl)amino-6-methoxy-2-(1-imidazolyl)quinazoline or
 4-(2-phenoxyethyl)amino-2-(1-imidazolyl)quinazoline,
 and pharmaceutically acceptable acid addition salts

formula


U.S. Patent No. 5,576,322 discloses compounds of the




wherein R1, R3, and R4, each of which may be the same or different from each other, may each represent a hydrogen atom, a halogen atom or a lower alkyl group or a lower alkoxy hydrogen atom, R2 is a halogen or cyan group R5 is a group represented by the formula:




wherein u is 3 or 4 and R61 represents a carboxyl group which may be protected or a heteromaryl group; or R5 is a group represented by the formula:



and R6 is a group represented by the formula



wherein X is hydrogen atom or a halogen atom or




or the pharmacologically acceptable salt thereof

Preferred compounds include:

2-(4-carboxypiperidino)-4-(3,4-methylene-dioxybenzyl) amino-6-chloroquinazoline- or a pharmaceutically acceptable salt thereof.

Sodium 2-(4-carboxypiperidino)-4-(3,4-methylene-dioxybenzyl) amino-6-chloroquinazoline.

WO 94/29277 discloses compounds of the formula



Formula (1)

or a pharmaceutically acceptable salt thereof, wherein

Ar is an optionally substituted aryl or heteroaryl ring selected from phenyl, naphthyl, pyridyl, pyrimidyl, pyridazinyl, pyrazinyl, imidazolyl, thienyl, oxazolyl, benzimidazolyl, benzoxazolyl, indolyl or thianaphthenyl,

X is CH or N;

R⁰ is NR¹R² or hydrogen; and


R¹ and R² are independently hydrogen or C₁₋₆alkyl.

Preferred compounds include:

- 3-amino-4-[4-(3-pyridyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(4-imidazolyl)]anilino-3-cyclobutene-1,2-dione,
- 3-methylamino-4-[3-(5-methyl-4-imidazolyl)]anilino-3-cyclobutene-1,2-dione,
- 3-dimethylamino-4-[3-(5-methyl-4-imidazolyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(3-methyl-4-pyridyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(2-oxazolyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(4-pyridyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(3-pyridyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(2-pyridyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(2-thienyl)]anilino-3-cyclobutene-1,2-dione,
- 3-amino-4-[3-(3-thienyl)]anilin]-3-cyclobutene-1,2-dione,

3-amino-4-[3-(2-thianaphthenyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(5-pyrimidyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(2-benzoxazolyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(2-benzimidazolyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(2-indolyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-(3-phenyl)anilino-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(2-hydroxyphenyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(2-methoxyphenyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(3-hydroxy-2-pyridyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[3-(2-imidazolyl)anilino]-3-cyclobutene-1,2-dione,
 3-amino-4-[6-(4-pyridyl)-2-pyridylamino]-3-cyclobutene-1,2-dione, or
 3-[3-(4-pyridyl)anilino]-3-cyclobutene-1,2-dione,
 or a pharmaceutically acceptable salt thereof.

WO 95/19978 discloses compounds of the formula




and salts and solvates thereof, in which:

R⁰ represents hydrogen, halogen or C₁₋₆ alkyl;

R¹ represents hydrogen, C₁₋₆alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, haloC₁₋₆alkyl, C₃₋₈cycloalkyl, C₃₋₈cycloalkylC₁₋₃alkyl, arylC₁₋₃alkyl or heteroarylC₁₋₃alkyl;

R² represents an optionally substituted monocyclic aromatic ring selected from benzene, thiophene, furan and pyridine or an optionally




substituted bicyclic ring attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen; and R³ represents hydrogen or C₁₋₃ alkyl, or R¹ and R³ together represent a 3- or 4- membered alkyl or alkenyl chain.

-50-

Preferred compounds include:

Cis-2,3,6,7,12,12a-hexahydro-2-(4-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;
 (5aR,12R,14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1'',2'':4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indole-5-1,4-dione;
 and physiologically acceptable salts and solvates thereof.

WO 96/28429 discloses compounds of the formula



wherein:

- R¹ is tert-butyl, or cyclopentyl;
- R³ is methyl, ethyl, or phenylmethyl;
- X is -CH₂-, -O-, or -NH-; and
- R⁶ is phenyl (or phenyl substituted by from one to three, the same or different, substituents selected from the group

-51-

consisting of lower-alkoxy, hydroxy, halogen, carboxylower-alkoxy, 4-morpholinyl-lower-alkoxy, 5-tetrazolyl-lower-alkoxy, dilower-alkylamino, trifluoromethyl, nitro, amino, lower-alkylsulfonylamino, dilower-alkylamino-lower-alkylphenyl carbonyloxy, and 1-imidazolyl); or when X is -CH₂- R⁶ is additionally 2-, 3-, or 4-pyridinyl, 1-pyrrolyl, 1-benzimidazolyl, 1,2,3,4-tetrahydro-2-isoquinolinyl, 1,2,3,4-tetrahydro-1-quinolinyl, hydroxy, 1-imidazolyl, 1-lower-alkyl-2,3,4, or 5-pyrrolyl, 1-pyrazolyl, 3-, 4-, or 5-isoxazolyl (or 3,4, or 5-isoxazolyl substituted on any available carbon atom thereof by lower-alkyl), 2-thienyl, or 3-thienyl; or a pharmaceutically acceptable acid-addition salt and/or hydrate thereof.

Preferred compounds include:


1-cyclopentyl-3-ethyl-6-(4-methoxyphenylmethyl)pyrazolo [3,4-d]pyrimidin-4-one,

1-cyclopentyl-3-ethyl-6-(4-hydroxyphenylmethyl)pyrazolo [3,4-d]pyrimidin-4-one,

1-cyclopentyl-3-ethyl-6-(phenylmethyl)pyrazolo[3,4-d] pyrimidin-4-one, and

1-cyclopentyl-3-ethyl-6-(4-aminophenylmethyl)pyrazolo [3,4-d]pyrimidin-4-one.

WO 96/28448 discloses compounds of the formula



wherein:

R¹ is tert-butyl, or cyclopentyl;

R³ is lower-alkyl, or phenyl-lower-alkyl; and

R⁶ is phenyl, or phenyl substituted by from one to three, the same or different, substituents selected from the group consisting of lower-alkoxy, lower-alkyl, hydroxy, 1-imidazolyl,

lower-alkenyloxy, dilower-alkylamino-lower-alkoxy, 4-morpholinyl-lower-alkoxy, lower-alkoxycarbonyl-lower-alkoxy, carboxylower-alkoxy, trifluoromethyl-, 1-piperidinyl-lower-alkoxy, 1-pyrrolidinyl-lower-alkoxy, nitro, halo, amino, $-(CH_2)_2O-$, lower-alkylsulfonylamino, lower-alkoxy-lower-alkoxy, lower-alkenyl, dilower-alkylamino, $-OCH(CH_3)CH_2-$, 4-morpholinylcarbonyl-lower-alkoxy, 4-thiomorpholinyl-lower-alkoxy, pyridinyl-lower-alkoxy, 1-lower-alkyl-3-hexahydroazepinyloxy, and 1-lower-alkyl-4-piperidinyl oxy; or a pharmaceutically acceptable acid-addition salt and/or hydrate thereof.

Preferred compounds include:

1- cyclopentyl-3-ethyl-6-(2-propoxyphenyl)pyrazolo[3,4-d]pyrimidin-4-one,


1-cyclopentyl-3-ethyl-6-[4-(1-imidazolyl)phenyl]pyrazolo[3,4-d]pyrimidin-4-one,

1-cyclopentyl-3-ethyl-6-[3-(2-(4-morpholinyl)ethoxy)phenyl]pyrazolo[3,4-d]pyrimidin-4-one,

1-cyclopentyl-3-ethyl-6-[2-ethoxy-4-(1-imidazolyl)phenyl]pyrazolo[3,4-d]pyrimidin-4-one, and

1-cyclopentyl-3-ethyl-6-[2-($CH_2=CHCH_2O$)phenyl]pyrazolo[3,4-d]pyrimidin-4-one.

WO 96/32003 discloses compounds of the formula



and salts and solvates thereof, in which:


R^0 represents hydrogen, halogen or C_{1-6} alkyl;

R^1 is selected from the group consisting of:

- (a) hydrogen;
- (b) C_{1-6} alkyl optionally substituted by one or more substituents selected from phenyl, halogen, $-CO_2R^a$ and $-NR^aR^b$;
- (c) C_{3-8} cycloalkyl;
- (d) phenyl; and
- (e) a 5- or 6-membered heterocyclic ring containing at least one heteroatom selected from oxygen, nitrogen and sulphur, and being optionally substituted by one or more C_{1-6} alkyl, and optionally linked to the nitrogen atom to which R^1 is attached via C_{1-6} alkyl;

R^2 is selected from the group consisting of:

- (f) C_{3-8} cycloalkyl;
- (g) phenyl optionally substituted by one or more substituents selected from $-OR^a$, $-NR^aR^b$, halogen, hydroxy, trifluoromethyl, cyano and nitro;
- (h) a 5- or 6-membered heterocyclic ring containing at least one heteroatom selected from oxygen, nitrogen and sulphur; and



- (i) a bicyclic ring attached to the rest of the molecule via one of the benzene ring carbon atoms and A is a 5- or 6-membered heterocyclic ring as defined in point (h); and

R^a and R^b independently represent hydrogen or C_{1-6} alkyl.

Preferred compounds include:

- Cis-2-benzyl-5-(3,4-methylenedioxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-benzyl-5-(3,4-methylenedioxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-5-(4-methoxyphenyl)-2-methyl-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-ethyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-ethyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-ethyl-5-(3,4-methylenedioxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-ethyl-5-(2-thienyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-5-(4-dimethylaminophenyl)-2-ethyl-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-9-methyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-9-bromo-2-butyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-9-fluoro-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-9-fluoro-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(3,4-methylenedioxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(3-chlorophenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(3-chlorophenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(4-chlorophenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(4-chlorophenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(4-fluorophenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;


Trans-2-butyl-5-(4-hydroxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(4-trifluoromethylphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(4-cyanophenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(4-cyanophenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(4-nitrophenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(4-nitrophenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(3-pyridyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b] indole-1,3(2H)-dione;
Cis-2-butyl-5-(3-thienyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(3-thienyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-butyl-5-(3-furyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-(3-furyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-cyclohexyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-cyclohexyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-cyclohexyl-9-fluoro-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-cyclohexyl-9-fluoro-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-benzyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido [3,4-b]indole-1,3(2H)-dione;
Cis-2-benzyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido [3,4-b]indole-1,3(2H)-dione;
Trans-2-benzyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6] pyrido [3,4-b]indole-1,3(2H)-dione;
(5R,11aR)-2-benzyl-5-(3,4-methylenedioxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-benzyl-5-(4-hydroxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo [1',5':1,6] pyrido [3,4-b]indole-1,3(2H)-dione;

-56-

Trans-2-(2-chloroethyl)-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-benzyl-5-cyclohexyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-benzyl-5-cyclohexyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-butyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-cyclohexyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-2-cyclohexyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-ethoxycarbonylmethyl-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-(4-methoxyphenyl)-2-[2-(2-pyridyl)-ethyl]-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-cyclopropyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-phenethyl-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-phenyl-2-(2-pyridylmethyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-phenyl-2-(4-pyridylmethyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-(4-methoxyphenyl)-2-(3-pyridylmethyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-(2-dimethylamino-ethyl)-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-(3-dimethylamino-propyl)-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-2-(2-morpholin-4-yl-ethyl)-5-phenyl-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-(4-methoxyphenyl)-2-[3-(4-methyl-piperazin-1-yl)-propyl]-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-(4-methoxyphenyl)-2-(2-pyrrolidin-1-yl-ethyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-(4-methoxyphenyl)-2-[2-(1-methyl-pyrrolidin-2-yl)-ethyl]-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Trans-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
Cis-5-(4-methoxyphenyl)-5,6,11,11a-tetrahydro-1H-imidazo[1',5':1,6]pyrido[3,4-b]indole-1,3(2H)-dione;
and pharmaceutically acceptable salts and solvates thereof.

-57-

WO 96/32379 discloses compounds of the formula



wherein

R^1 is hydrogen, halogen, nitro, carboxy, protected carboxy, acyl, cyano, hydroxyimino(lower)alkyl, lower alkenyl optionally substituted with oxo, or lower alkyl optionally substituted with protected carboxy, carboxy or hydroxy;

R^2 is hydrogen, halogen, lower alkenyl, acyl, or lower alkyl optionally substituted with protected carboxy, carboxy, lower alkoxy or hydroxy;

R^3 is lower alkenyl or lower alkyl, both of which are optionally substituted with one or more substituent(s) selected from the group consisting of

(1) oxo,

(2) aryl optionally substituted with one or more substituent(s) selected from the group consisting of halogen, aryl, lower alkoxy, lower alkylendioxy, cyano, nitro, carboxy, protected carboxy, acyl, and amino optionally substituted with acyl or protected carboxy, and


(3) a heterocyclic group optionally substituted

with halogen; and

R^4 is carboxy, protected carboxy, acyl, cyano, halogen, a heterocyclic group, amino optionally substituted with acyl or protected carboxy, or lower alkyl

optionally substituted with protected carboxy, carboxy or acyl;
 in addition to their significances above,
 R^1 and R^2 , together with the carbon atoms to which they are attached, represent a 4- to 7-membered carbocyclic ring optionally substituted with oxo, or its pharmaceutically acceptable salt.

WO 97/03070 discloses compounds of the formula



wherein R^1 is a hydrogen atom or a halogen atom;

R^2 is a phenyl-lower alkyl group;

R^3 is a heterocyclic group selected from the group consisting of an indolyl group, indolinyl group, 1H-indazolyl group, 2(1H)-quinolinonyl group, 3,4-dihydro-2(1H)-quinolinonyl group and 3,4-dihydro-1,4(2H)-benzoxazinyl group, said heterocyclic group may have 1 to 3 substituents selected from the group consisting of:

a group of the formula $-B-R^4$, (B is a lower alkylene group; R^4 is a 5- to 11-membered saturated or unsaturated heterocyclic group of single ring or binary ring, having 1 to 4 hetero atoms selected from the group consisting of a nitrogen atom, oxygen atom and sulfur atom, (said heterocyclic group may have 1 to 3 substituents selected from the group consisting of a halogen atom, a lower alkyl group, a lower alkoxy group and

-59-

oxo group) or a group of the formula $-NR^5R^6$ (R^5 and R^6 are each the same or different, and a hydrogen atom, a lower alkyl group, a cycloalkyl group, a pyridyl-carbonyl group, an isoxazolylcarbonyl group which may have 1 to 3 lower alkyl groups as the substituents, a pyrrolylcarbonyl group or an amino-substituted lower alkyl group which may have a lower alkyl group as the substituent; further R^5 and R^6 may form 5- to 6-membered saturated heterocyclic group by combining to each other, together with the adjacent nitrogen atom being bonded thereto, further with or without other nitrogen atom or oxygen atom; said heterocyclic group may have 1 to 3 substituents selected from the group consisting of a hydroxy group and a phenyl group)); a lower alkenyl group; a lower alkoxycarbonyl group; a phenoxy-lower alkyl group which may have cyano group as the substituents; a halogen-substituted lower alkyl group; and a lower alkoxycarbonyl-substituted lower alkyl group;

A is a lower alkylene group; and

n is 0 or 1.

Preferred compounds include:

1-Benzyl-6-chloro-2-{1-[3-(imidazol-1-yl)propyl]indol-5-ylaminocarbonyl}benzimidazole.

1-Benzyl-6-chloro-2-{1-[3-(N-cyclohexyl-N-methylamino)propyl]indol-5-ylaminocarbonyl}-benzimidazole.

-60-

1-Benzyl-6-chloro-2-{1-[3-(pyrazol-1-yl)propyl]indol-5-ylaminocarbonyl}benzimidazole.


1-Benzyl-6-chloro-2-{1-[3-(1,2,4-triazol-1-yl)propyl]indol-5-ylaminocarbonyl}benzimidazole.

1-Benzyl-6-chloro-2-{1-[3-(3,5-dimethylisoxazol-4-ylcarbonylamino)propyl]indol-5-ylaminocarbonyl}benzimidazole.

1-Benzyl-6-chloro-2-{1-[3-(4-phenyl-4-hydroxypiperidin-1-yl)propyl]indol-5-ylaminocarbonyl}benzimidazole.

1-Benzyl-6-chloro-2-{4-[3-(pyridin-2-ylcarbonylamino)propyl]-3,4-dihydro-1,4(2H)-benzoxazin-7-ylaminocarbonyl}benzimidazole.

WO 97/03675 discloses compounds of the formula




and salts and solvates (e.g. hydrates) thereof, in which:

R⁰ represents hydrogen, halogen or C₁₋₆ alkyl;

R¹ represents hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, haloC₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkylC₁₋₃ alkyl, arylC₁₋₃ alkyl or heteroarylC₁₋₃ alkyl;

R² represents an optionally substituted monocyclic aromatic ring selected from benzene, thiophene, furan and pyridine or an optionally substituted bicyclic



ring attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen; and

R³ represents hydrogen or C₁₋₃ alkyl, or R¹ and R³ together represent a 3- or 4- membered alkyl or alkenyl chain;

-61-

for the manufacture of a medicament for the curative or prophylactic treatment of erectile dysfunction in a male animal, including man.

Preferred compounds include:

Cis-2,3,6,7,12,12a-hexahydro-2-(4-pyridylmethyl)-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(2,3-dihydrobenzo[b]furan-5-yl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-6-(5-bromo-2-thienyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(5aR,12R,14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1",2":4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indole-5-1,4-dione;


Cis-2,3,6,7,12,12a-hexahydro-2-cyclopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

(3S,6R,12aR)-2,3,6,7,12,12a-hexahydro-3-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;

and physiologically acceptable salts and solvates (e.g. hydrates) thereof.

-62-

WO 97/03985 discloses compounds of the formula




and solvates thereof, in which:

R^0 represents hydrogen, halogen or C_{1-6} alkyl;

R^1 represents hydrogen or C_{1-6} alkyl;

R^2 represents the bicyclic ring



which may be optionally substituted by one or more groups selected from halogen and C_{1-3} alkyl;

and

R^3 represents hydrogen or C_{1-3} alkyl.

Preferred compounds include:

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(3S, 6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-3-methyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;


(3S, 6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2,3-dimethyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

(6R, 12aR)-2,3,6,7,12,12a-Hexahydro-6-(5-benzofuranyl)-2-isopropyl-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione;

and physiologically acceptable solvates thereof.

-63-

WO 97/43287 discloses compounds of the formula



wherein

R° represents -hydrogen or -halogen;

R¹ is selected from the group consisting of:

-hydrogen,

-NO₂,

-trifluoromethyl,

-trifluoromethoxy,

-halogen,

-cyano,

a 5- or 6- membered heterocyclic group containing at least one heteroatom selected from oxygen, nitrogen and sulphur (optionally substituted by - C(=O)OR* or C₁-₄alkyl),

-C₁-₄alkyl optionally substituted by -OR*,

-C₁-₃alkoxy,

-C(=O)R*,

-O-C(=O)R*,

-C(=O)OR*,

-C₁-₄alkylene C(=O)OR*,

-O-C₁-₄alkylene -C(=O)OR*,

-C₁-₄alkylene-O-C₁-₄alkylene-C(=O)OR*,

-C(=O)NR*SO₂R*,

-C(=O)C₁-₄alkylene Het, wherein Het represents 5- or 6-membered heterocyclic group as defined above,

-C₁-₄alkylene NR*R*,

-C₂-₄alkenylene NR*R*,

-C(=O)NR*R*,

-C(=O)NR*R*,

-C(=O)NR*C₁-₄alkylene OR*

-C(=O)NR*C₁-₄alkylene Het, where in Het represents a 5- or 6-membered

-64-

heterocyclic group as defined above,

-OR^a

-OC₂₋₄alkylene NR^aR^b,

-OC₁₋₄alkylene-CH(OR^a)CH₂ NR^aR^b,

-O-C₁₋₄alkylene Het, wherein Het represents a 5- or 6- membered heterocyclic group as defined above,

-O-C₂₋₄alkylene-OR^a,

-O-C₂₋₄alkylene-NR^a-C(=O)-OR^b,

-NR^aR^b,

-NR^aC₁₋₄alkyleneNR^aR^b,

-NR^aC(=O)R^b,

-NR^aC(=O)NR^aR^b,

-N(SO₂C₁₋₄alkyl)₂,

-NR^a(SO₂C₁₋₄alkyl),

-SO₂NR^aR^b, and

-OSO₂trifluoromethyl;

R² is selected from the group consisting of:

-hydrogen,

-halogen,

-OR^a,

-C₁₋₆ alkyl,

-NO₂, and

-NR^aR^b,

or R¹ and R², together form a 3- or 4- membered alkylene or alkenylene chain, optionally containing at least one heteroatom;

R³ is selected from the group consisting of:

-hydrogen,

-halogen,

-NO₂,

-trifluoromethoxy,

-C₁₋₆alkyl, and

-C(=O)OR^a;

R⁴ is hydrogen,

or R³ and R⁴ together form a 3- or 4- membered alkylene or alkenylene chain, optionally containing at least one heteroatom;

R^a and R^b, which may be the same or different, are independently selected from hydrogen and C₁₋₆alkyl;

R^c represents phenyl or C₄₋₆cycloalkyl, which phenyl or C₄₋₆cycloalkyl can be optionally substituted by one or more halogen atoms, one or more -C(=O)OR^a or one or more -OR^a;


-65-

n is an integer selected from 1, 2 and 3;


m is an integer selected from 1 and 2;

and pharmaceutically acceptable salts and solvates thereof.

U.S. Patent No. 5,393,755 discloses compounds of the formula



or



wherein

J is oxygen or sulfur,

R¹ is hydrogen, alkyl or alkyl substituted with aryl or hydroxy;

R² is hydrogen, aryl, heteroaryl, cycloalkyl, alkyl or alkyl substituted with aryl, heteroaryl, hydroxy, alkoxy, amino, monoalkyl amino or dialkylamino, or $-(CH_2)_mTCOR^{20}$ wherein m is an integer from 1 to 6, T is oxygen or $-NH-$ and R²⁰ is hydrogen, aryl, heteroaryl, alkyl or alkyl substituted with aryl or heteroaryl;

R³ is hydrogen, halo, trifluoromethyl, alkoxy, alkylthio, alkyl, cycloalkyl, aryl, aminosulfonyl, amino, monoalkylamino, dialkylamino, hydroxyalkylamino, aminoalkylamino, carboxy, alkoxy-carbonyl or aminocarbonyl or alkyl substituted with aryl, hydroxy, alkoxy, amino, monoalkylamino or dialkylamino;

R⁴, R⁵, R⁶ and R⁷ independently represent hydrogen, alkyl, cycloalkyl or aryl; or (R⁴ and R⁵) or (R⁶ and R⁷) or (R⁵ and R⁶) can complete a saturated ring of 5- to 7-carbon atoms, or (R⁴ and R⁵) taken together and (R⁶ and R⁷) taken together, each complete a saturated ring of 5- to 7-carbon atoms, wherein each ring optionally can contain a sulfur or oxygen atom and whose carbon atoms may be optionally substituted with one or more of the following: alkenyl, alkynyl, hydroxy, carboxy, alkoxy-carbonyl, alkyl or alkyl substituted with hydroxy, carboxy or alkoxy-carbonyl; or such saturated ring can have two adjacent carbon atoms which are shared with an adjoining aryl ring; and

n is zero or one.

Preferred compounds include:

cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(phenylmethyl)cyclopenta[4,5]imidazo[2,1-b]purin-4-one;
 7,8-Dihydro-5-methyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 cis-6a,7,8,9,10,10a-Hexahydro-5-methyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
 5,7,8,9-Tetrahydro-5-methyl-3-(phenylmethyl)-pyrimido[2,1-b]purin-4(3H)-one;
 7,8-Dihydro-8-phenyl-5-methyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 5', 7'-Dihydro-5'-methyl-3'-(phenylmethyl)spiro[cyclohexane-1,8'-(8H)imidazo[2,1-b]purin]-4'(3H)-one;
 cis-5,6a,11,11a-Tetrahydro-5-methyl-3-(phenylmethyl)indeno[1',2':4,5]imidazo[2,1-b]purin-4(3H)-one;
 5',7'-Dihydro-2',5'-dimethyl-3'-(phenylmethyl)spiro[cyclohexane-1,7'-(8'H)-imidazo[2,1-b]purin]-4'(3'H)-one;
 7,8-Dihydro-2,5,7,7,8(R,S)-pentamethyl-3H-imidazo[2,1-b]purin-4(5H)-one;
 cis-5,6a,7,11b-Tetrahydro-5-methyl-3-(phenylmethyl)indeno[2',1':4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4-(3H)-one;
 5'-Methyl-3'-(phenylmethyl)spiro[cyclopentane-1,7'-(8'H)-(3'H)imidazo[2,1-b]purin]-4-(5'H)-one;
 7,8-Dihydro-2,5,7,7-tetramethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-7(R)-phenyl-2,5-dimethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-3,7(R)-bis(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 (±)-7,8-Dihydro-2,5-dimethyl-7-ethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 6a(S)-7,8,9,10,10a(R)-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
 6a(R)-7,8,9,10,10a(S)-hexahydro-2,5-dimethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-7(R)-isopropyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5,7(R)-trimethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 cis-7,7a,8,9,10,10a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-3H-cyclopenta[5,6]pyrimido[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-7(S)-(1-methylpropyl)-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-7(R)-(2-methylpropyl)-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-7(R,S)-(methoxycarbonyl)-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-7(R,S)-(1-propyl)-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-7(S)-(1-methylethyl)-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5,7,7,8(R,S)-pentamethyl-3H-imidazo[2,1-b]purin-4(5H)-one;
 5,7,8,9-Tetrahydro-2,5,7,9(R,S)-pentamethyl-3-(phenylmethyl)-pyrimido[2,1-b]purin-4(3H)-one;
 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 5,6a(S),7,8,9,9a(R)-Hexahydro-2,5-dimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-6a,7,8,9,10,10a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;

5',7'-Dihydro-2',5'-dimethyl-3'-(phenylmethyl)spiro[cyclohexane-1,8-(8H)-imidazo[2,1-b]purin]-4-(3'H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2,5-dimethyl-3-(phenylmethyl)cyclohept[6,7]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-ethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-6a,7,8,9,10,10a-Hexahydro-5-methyl-2-ethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-ethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-phenyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-6a,7,8,9,10,10a-Hexahydro-5-methyl-2-phenyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methylcyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2,5-dimethylcyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a(R), 7,8,9,9a(S)-Hexahydro-2,5-dimethylcyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 2',5'-dimethyl-spiro[cyclopentane-1,7'-(8'H)-(3'H)-imidazo[2,1-b]purin]-4'(5'H)-one;
 7,8-Dihydro-2,5-dimethyl-7(R)-(1-methylethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5,7,7-tetramethyl-3H-imidazo[2,1-b]purin-4(5H)-one;
 7,8-Dihydro-2,5-dimethyl-7(S)-(1-methylethyl)-3H-imidazo[2,1-b]purin-4(5H)-one;
 6a(R),7,8,9,10,10a(S)-Hexahydro-2,5-dimethyl-3H-benzimidazo[2,1-b]purin-4(5H)-one;
 5',7'-Dihydro-2',5'-dimethylspiro[cyclohexane-1,7-(8'H)-imidazo[2,1-b]purin]-4'(3'H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(phenylmethyl)cyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-thione;
 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-thione;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(4-chlorophenylmethyl)cyclopenta[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(cyclohexylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-3-(2-naphthylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(4-bromophenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 5,6a(R)-7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(4-methoxyphenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2,3,5-trimethylcyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2-(hydroxymethyl)-5-methyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-5,6a,7,8,9,9a-Hexahydro-2-methylthio-5-methyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;
 cis-3,4,5,6a,7,8,9,9a-Octahydro-5-methyl-4-oxo-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-2-carboxylic acid;
 cis-3,4,5,6a,7,8,9,9a-Octahydro-5-methyl-4-oxo-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-2-

carboxylic acid, methyl ester;
 cis-5,6a,7,8,9,9a-Hexahydro-2-bromo-5-methyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 cis-5,6a,7,8,9,9a-Hexahydro-2-(methylaminosulfonyl)-5-methyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 cis-1-Cyclopentyl-5,6a,7,8,9,9a-hexahydro-5-methylcyclopent[4,5]imidazo[2,1-b]purin-4(1H)one;
 cis-5,6a,7,8,9,9a-Hexahydro-3,5-bis-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 cis-6a,7,8,9,10,10a-Hexahydro-3,5-bis-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)one;
 cis-3-Cyclopentyl-5,6a,7,8,9,9a-hexahydro-5-methylcyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 5'-Methyl-3'-(phenylmethyl)spiro[cyclopentane-1,7-(8'H)-(3'H)imidazo[2,1-b]purin]-4(5H)one;
 2',5'-Dimethyl-3'-(phenylmethyl)-spiro[cyclopentane-1,7-(8'H)-(3'H)imidazo[2,1-b]purin]-4(5'H)one;
 cis-5,6a,(R),7,8,9,9a(S)-Hexahydro-5-methyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 cis-3-Cyclopentyl-5,6a,7,8,9,9a-Hexahydro-2,5-dimethylcyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 5'-Methyl-2'-trifluoromethyl-3'-(phenylmethyl)spiro[cyclopentane-1,7'-(8'H)-(3'H)imidazo[2,1-b]purin]-4(5'H)one;
 7,8-Dihydro-5,7,7-trimethyl-2-trifluoromethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(5H)one;
 (+/-)-cis-5,6a,7,8,9,9a-Hexahydro-5-methyl-2-trifluoromethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 (+/-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3-(phenylmethyl)-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)one;
 (+)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3-phenylmethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)one;
 (-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3-phenylmethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)one;
 (+/-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)one;
 (+)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)one;
 (-)-6a,7,8,9,9a,10,11,11a-Octahydro-2,5-dimethyl-3H-pentaleno[6a',1':4,5]imidazo[2,1-b]purin-4(5H)one;
 6a,7,8,9,10,10a,11,12,13,13a-Decahydro-2,5-dimethyl-3-(phenylmethyl)naph[1,8a-d]imidazo[2,1-b]purin-4(5H)one;
 7(R)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(3H)one;
 7(R)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3H-imidazo[2,1-b]purin-4(5H)one;
 7(S)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3-(phenylmethyl)-3H-imidazo[2,1-b]purin-4(3H)one;
 7(S)-Cyclohexyl-7,8-dihydro-2,5-dimethyl-3H-imidazo[2,1-b]purin-4(5H)one;
 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(trimethylacetoxymethyl)-cyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;
 5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-(4-pyridylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)one;

-69-

5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-[2-(1-morpholinyl)ethyl]cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;

5,6a(R),7,8,9,9a(S)-Hexahydro-2,5-dimethyl-3-[acetoxymethyl]cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;

5,6a,7,8,9,9a-Hexahydro-2,5,6a-trimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;

5,6a(R),7(S),8,9,9a-Hexahydro-2,5,6a-trimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;


5,6a(S),7(R),8,9,9a-Hexahydro-2,5,6a-trimethyl-3-(phenylmethyl)cyclopent[4,5]imidazo[2,1-b]purin-4(3H)-one;

cis-6a,7,8,9,10,10a-Hexahydro-2,5,7-trimethyl-3-(phenylmethyl)-3H-benzimidazo[2,1-b]purin-4(5H)-one;

cis-5,6a,7,8,9,9a-Hexahydro-2,5,6a-trimethylcyclopent[4,5]imidazo[2,1-b]purin-4(3H); or

cis-6a,7,8,9,10,10a-Hexahydro-2,5,7-trimethyl-3H-benzimidazo[2,1-b]purin-4(5H)-one].

U.S. Patent No. 5,439,895 discloses compounds of the formula



wherein R¹ is hydrogen or C1-4 alkyl;

Y is C1-6 alkylene;

A is —O—R⁰ or —S(O)_p—R⁰,

in which R⁰ is C1-4 alkyl-hydroxy;

p is 0-2;

Z is single bond, methylene, ethylene, vinylene or ethynylene;

CyB is

- (1) 7-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one, two or three nitrogen atoms,
- (2) 6-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, two or three nitrogen atoms,
- (3) 6-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atom, one nitrogen atom,
- (4) 4- or 5-membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one, two or three nitrogen atoms, or
- (5) 4-7 membered, unsaturated or partially saturated, monocyclic hetero ring containing as hetero atoms, one or two oxygen atoms, or one or two sulfur atoms;

R³ is hydrogen, C1-4 alkyl, C1-4 alkoxy, halogen or trifluoromethyl;

R⁴ is (1) hydrogen, (2) C1-4 alkyl, (3) C1-4 alkoxy, (4) —COOR⁵, in which R⁵ is hydrogen or C1-4 alkyl, (5) —NR⁹R¹⁰, in which R⁹ is hydrogen, C1-4 alkyl or phenyl(C1-4 alkyl) and R¹⁰ is hydrogen or C1-4 alkyl, (6) —NHCOR¹¹, in which R¹¹ is C1-4 alkyl, (7) —NHSO₂R¹¹, in which R¹¹ is as hereinbefore defined, (8) SO₂NR⁹R¹⁰, in which R⁹ and R¹⁰ are as hereinbefore defined, (9) —OCOR¹¹, in which R¹¹ is as hereinbefore defined, (10) halogen, (11) trifluoromethyl, (12) hydroxy, (13) nitro,

-70-

(14) cyano, (15) $-\text{SO}_2\text{N}=\text{CHNR}^{12}\text{R}^{13}$ in which R^{12} is hydrogen or C1-4 alkyl and R^{13} is C1-4 alkyl, (16) $-\text{CONR}^{14}\text{R}^{15}$ in which R^{14} is hydrogen or C1-4 alkyl and R^{15} is C1-4 alkyl or phenyl(C1-4 alkyl), (17) C1-4 alkylthio, (18) C1-4 alkylsulfinyl, (19) C1-4 alkylsulfonyl, (20) ethynyl, (21) hydroxymethyl, (22) tri(C1-4 alkyl)silylethynyl or (23) aceryl; and m and n independently are 1 or 2; with the proviso that

(1) a CyB ring does not bond to Z through a nitrogen atom in the CyB ring when Z is vinylene or ethynylene;


or pharmaceutically acceptable acid addition salts thereof, pharmaceutically acceptable salts thereof, or hydrates thereof.

Preferred compounds include:

4-[2-(2-hydroxyethoxy)ethyl]amino-6-acetyl-2-(1-imidazolyl)quinazoline,
 2-(1-imidazolyl)-4-[2-(2-hydroxyethoxy)ethyl]amino-6-ethynylquinazoline,
 2-(1-imidazolyl)-4-[2-(2-hydroxyethoxy)ethyl]amino-6-(2-triisopropylsilylethynyl)quinazoline,
 4-[2-(2-hydroxyethoxy)ethyl]amino-6-hydroxymethyl-2-(1-imidazolyl)quinazoline,
 4-(2-(2-hydroxyethoxy)ethyl)amino-6-methylsulfinyl-2-(1-imidazolyl)quinazoline,
 6-chloro-4-(2-(2-hydroxyethoxy)ethyl)amino-2-(1-imidazolyl)quinazoline,
 4-[2-(2-hydroxyethoxy)ethyl]amino-6-methoxycarbonyl-2-(1-imidazolyl)quinazoline,
 4-(2-(2-hydroxyethoxy)ethyl)amino-6-methylthio-2-(1-imidazolyl)quinazoline,
 4-(2-(2-hydroxyethoxy)ethyl)amino-6-iodo-2-(1-imidazolyl)quinazoline,
 4-(2-(2-hydroxyethoxy)ethyl)amino-2-(1-imidazolyl)-5,6,7,8-tetrahydroquinazoline or
 6-methoxy-4-(2-(2-hydroxyethoxy)ethyl)amino-2-(1-imidazolyl)quinazoline,
 and pharmaceutically acceptable acid addition salts thereof, pharmaceutically acceptable salts thereof, or hydrates thereof.

-71-

U.S. Patent No. 5,488,055 discloses compounds of the formula



wherein:

R¹ is lower-alkyl, phenyl-lower-alkyl, or cycloalkyl;

R² is hydrogen, or lower-alkyl;

R³ is hydrogen, lower-alkyl, or hydroxylower-alkyl;


R⁴ is cycloalkyl or cycloalkyl substituted by from one to two, the same or different, substituents selected from the group consisting of lower-alkoxycarbonyl, carboxy, lower-alkylthio-lower-alkoxycarbonyl, hydroxylower-alkyl, hydroxy, oxo, lower-alkoxy, lower-alkyl, and halogen; and

R⁵ is from one to three, the same or different, substituents selected from the group consisting of hydrogen, lower-alkoxy, hydroxy, dilower-alkylamino-lower-alkoxy, carboxylower-alkoxy, lower-alkoxycarbonyl-lower-alkoxy, nitro, polyhydroxylower-alkoxy, amino, epoxy-lower-alkoxy, carboxy, lower-alkanoylamino, lower-alkoxycarbonyl, pyridinyl, 4-morpholinyl-lower-alkoxy, lower-alkylsulfonyl, cyano, 1-imidazolyl, halogen, dilower-alkylaminosulfonyl, oxadiazolyl (or oxadiazolyl substituted on any available carbon atom thereof by lower-alkyl), lower-alkylsulfinyl, 1-pyrazolyl (or 1-pyrazolyl substituted on any available carbon atom thereof by lower-alkyl), trifluoromethylsulfonyl, lower-alkenyl, lower-alkyl, and lower-alkynyl; or a pharmaceutically acceptable acid-addition salt and/or hydrate and/or solvate thereof, or, where applicable, a stereoisomer or a racemic mixture thereof.

Preferred compounds include

- 1-ethyl-6-nitro-N-[S(+)-1-(cyclohexyl)ethyl]-1H-pyrazolo [3,4-b]quinolin-4-amine,
- 1-ethyl-6-nitro-N-[cyclohexylmethyl]-1H-pyrazolo [3,4-b]quinolin-4-amine,
- 1-ethyl-6-cyano-N-[S(+)-1-(cyclohexyl)ethyl]-1H-pyrazolo [3,4-b]quinolin-4-amine,
- 1-ethyl-6-bromo-N-[S(+)-1-(cyclohexyl)ethyl]-1H-pyrazolo [3,4-b]quinolin-4-amine, and
- 1-ethyl-6-(1-pyrazolyl)-N-[S(+)-1-(cyclohexyl)ethyl]-1H-pyrazolo [3,4-b]quinolin-4-amine.

U.S. Patent No. 5,525,064 discloses compounds of the formula



wherein A is a bond, C₁₋₄ alkylene or C₁₋₄ oxyalkylene;
Y is a bond, C₁₋₄ alkylene, C₁₋₄ alkyleneoxy, C₁₋₄ alkox-
yphenylene or phenyl(C₁₋₄)alkylene;

Z is a bond or vinylene;

R¹ is a heterocyclic ring selected from the group consist-
ing of pyrrole, pyridine, azepine, imidazole, pyrazole,
pyrimidine, pyrazine, pyridazine, benzimidazole,
quinoline, isoquinoline and partially or fully saturated
rings thereof;

R² is

(i) a heterocyclic ring selected from the group consist-
ing of pyrrole, pyridine, azepine, imidazole, pyra-
zole, pyrimidine, pyrazine, pyridazine, benzimida-
zole, quinoline, isoquinoline, furan, pyran, dioxole,
dioxine, benzofuran, benzopyran, benzodioxole,
benzodioxine, thiophene, thioine, benzothiophene,
benzathione and partially or fully saturated rings
thereof,

(ii) C₄₋₁₅ carbocyclic ring,

(iii) C₁₋₄ alkoxy,

(iv) hydroxy(C₁₋₄ alkoxy), or

(v) hydroxy;

with the proviso that:

when R¹ is pyridine or pyridine substituted by one or
two of C₁₋₄ alkyl,

C₁₋₄ alkoxy, halogen, trifluoromethyl or nitro then R²
is a member selected only from the group con-
sisting of benzodioxole or benzodioxole substi-
tuted by one or two of C₁₋₄ alkyl, C₁₋₄ alkoxy,
halogen, trifluoromethyl, nitro or a group of the
formula:



wherein R¹⁰ is hydrogen or C₁₋₄ alkyl, and
hydroxy(C₁₋₄ alkoxy);

R³ is

(i) a heterocyclic ring selected from the group consist-
ing of pyrrole, pyridine, azepine, imidazole, pyra-
zole, pyrimidine, pyrazine, pyridazine, benzimida-
zole, quinoline, isoquinoline, furan, pyran,
benzofuran, benzopyran, thiophene, thioine, ben-
zothiophene, benzathione, thiazole, isothiazole, thia-
zine, benzothiazole, benzoisothiazole, benzothiaz-
ine and partially or fully saturated rings thereof,

(ii) C₄₋₁₅ carbocyclic ring,

(iii) a group of formula:



wherein X is halogen, or

(iv) hydrogen,

l is 1 or 2,

with the proviso that:

the ring represented by R¹ may be substituted by one or
two of C₁₋₄ alkyl, C₁₋₄ alkoxy, halogen, trifluorom-
ethyl or nitro;

the ring represented by R² may be substituted by one or
two of C₁₋₄ alkyl, C₁₋₄ alkoxy, halogen, trifluorom-
ethyl, nitro or a group of the formula:

-73-

-COOR¹⁰

wherein R¹⁰ is hydrogen or C₁₋₄ alkyl, and the ring represented by R³ may be substituted by one or two of C₁₋₄ alkyl, C₁₋₄ alkoxy, halogen, trifluoromethyl, nitro, cyano, ethynyl or a group of the formula:

-SONR⁷R⁸

wherein R⁷ and R⁸ are independently hydrogen or C₁₋₄ alkyl, and with the proviso that

R² is not hydroxy when Y is a bond; and

R¹ is not bonded through its nitrogen atom when Z is vinylene,

or pharmaceutically acceptable acid addition salts thereof or pharmaceutically acceptable salts thereof.


Preferred compounds include

2-(1-Imidazolyl)-4-[2-(2-hydroxyethoxy)ethyl]amino-5-(3-methoxyphenyl)methylpyrimidine,
 2-(1-Imidazolyl)-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-4-(2-methoxyethyl)aminopyrimidine,
 2-(1-Imidazolyl)-5-ethyl-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-phenylmethyl-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-methyl-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5,6-dimethyl-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-(3-methoxyphenyl)methyl-4-(2-methoxyethyl)aminopyrimidine,
 2-(1-Imidazolyl)-5-(4-methoxyphenyl)methyl-4-[2-(2-hydroxyethoxy)ethyl]aminopyrimidine,
 2-(1-Imidazolyl)-5-(4-methoxyphenyl)methyl-4-(2-methoxyethyl)aminopyrimidine or
 2-(1-Imidazolyl)-5-(4-methoxyphenyl)methyl-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-phenoxymethyl-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-(1-Imidazolyl)methyl-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-(1-chlorovinyl)-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thiazolyl)-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-[2-(2-hydroxyethoxy)ethyl]aminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(1-naphthyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(4-methoxyphenyl)methylaminopyrimidine,

2-(1-Imidazolyl)-5-(2-thienyl)-4-(3-methoxyphenyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(2-furyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(2-thienyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(3-pyridyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(2-methoxyethyl)aminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-phenylmethoxyaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(4-chlorophenyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(3-chlorophenyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(4-methylphenyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(4-methoxyphenyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(5-methyl-2-thienyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-[4-(1-imidazolyl)phenyl]methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(3-pyridyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(3-furyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(3-pyridyl)-4-phenylmethylaminopyrimidine,
 2-(1-Imidazolyl)-5-(4-chlorophenyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(benzimidazol-1-yl)-5-(2-thienyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(4-ethoxycarbonylphenyl)methylaminopyrimidine,
 2-(1-Imidazolyl)-5-(2-naphthyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(3-Pyridyl)-5-(2-thienyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-[2-(3-Pyridyl)vinyl]-5-(2-thienyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine,
 2-(2-Methyl-1-Imidazolyl)-5-(2-thienyl)-4-(1,3-dioxaindan-5-yl)methylaminopyrimidine or
 2-(1-Imidazolyl)-5-(2-thienyl)-4-(benzimidazol-5-yl)methylaminopyrimidine.

-75-


European published patent application No. 0728759
discloses compounds of the formula



wherein



is a heterocycle selected from



n is 0, 1 or 2;

Y is single bond or C1-6 alkylene;

Z is single bond, C1-2 alkylene or vinylene;

E is

(i) 4-15 membered, unsaturated, partially saturated or fully saturated, mono or bicyclic hetero ring containing one or two hetero atoms, chosen from nitrogen, oxygen and sulfur, not more than one hetero atom being sulfur,

(ii) 4-15 membered, unsaturated or partially saturated, mono or bicyclic carbocyclic ring, or

(iii) $-OR^4$; in which R^4 is hydrogen atom, C1-4 alkyl or C1-4 alkyl substituted by a hydroxy group;

Cyc is 5-7 membered, unsaturated, partially saturated or fully saturated, monocyclic hetero ring containing one or two nitrogen atoms or 5-7 membered, unsaturated or partially saturated, monocyclic carbocyclic ring;

R^1 is hydrogen atom or C1-4 alkyl;

R^2 is hydrogen atom, C1-4 alkyl, C1-4 alkoxy or halogen atom;


R^3 is hydrogen atom, C1-4 alkyl, C1-4 alkoxy or $-COOR^5$; in which R^5 is hydrogen atom or C1-4 alkyl;

with the proviso that

(1) a Cyc ring does not bond to Z through a nitrogen atom in the Cyc ring where Z is vinylene and that

(2) Y is not a single bond, when E is $-OR^4$; or a pharmaceutically acceptable acid addition salt, pharmaceutically acceptable salt or hydrate thereof.

U.S. Patent No. 5,541,187 discloses compounds of the formula



wherein:

R^1 is hydrogen, alkyl, cycloalkyl, cycloalkyl substituted by alkyl or hydroxyl, 2- or 3-tetrahydrofuranyl, 3-tetrahydrothienyl, 1,1-dioxide, cycloalkyl-alkyl, carboxy-alkyl, carbo-lower-alkoxy-alkyl, dialkylaminoalkyl,

phenyl-lower-alkyl, phenyl-lower-alkyl in which the phenyl ring is substituted in the 2, 3, or 4-position by one or two substituents, the same or different, selected from the group consisting of amino, halogen, alkyl, carboxyl, carbo-lower-alkoxy, carbamoyl, NHSO_2 - (quinolinyl), nitro and cyano;

R^3 is hydrogen, lower-alkyl, phenyl-lower-alkyl, lower-alkoxyphenyl-lower-alkyl, dilower-alkoxy-phenyl-lower-alkyl, pyridyl-lower-alkyl, cycloalkyl-lower-alkyl, phenylamino, dialkylamino, halogen, trifluoromethyl, lower-alkylthio, cyano or nitro; and

R^2 is a five or six membered heterocyclic ring containing from one to two nitrogen atoms, substituted—or unsubstituted—at any available carbon atom by one or two substituents, the same or different, selected from the group consisting of lower-alkyl, halogen, lower-alkoxy, cycloalkyloxy, 4-morpholinyl, lower-alkoxy-lower-alkoxy, hydroxy, imidazolyl, oxo and 4-morpholinyl-lower-alkoxy; or at any available nitrogen atom by lower-alkyl, lower-alkanoyl, or trifluoroacetyl; or a pharmaceutically acceptable acid-addition salt thereof.

Preferred compounds include:

1-Cyclopentyl-3-methyl-6-(4-pyridyl)pyrazolo[3,4-d]pyrimidin-4-one,


1-Cyclopentyl-3-ethyl-6-(3-ethoxy-4-pyridyl)pyrazolo[3,4-d]pyrimidin-4-one,

1-Cyclopentyl-3-ethyl-6-(3-methoxy-4-pyridyl)pyrazolo[3,4-d]pyrimidin-4-one,

1-Cyclopentyl-3-trifluoromethyl-6-(3-ethoxy-4-pyridyl)pyrazolo[3,4-d]pyrimidin-4-one,

1-Cyclopentyl-3-ethyl-6-(2-(1-imidazolyl)-4-pyridyl)pyrazolo[3,4-d]pyrimidin-4-one,


U.S. Patent No. 5,721,238 discloses compounds of the formula



(I)

in which

A represents oxiranyl, which is optionally substituted by straight-chain or branched alkyl having up to 8 carbon atoms, which in turn can be substituted by phenyl, or represents a radical of the formula



wherein

R¹ denotes hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,

R² denotes straight-chain or branched alkyl having up to 8 carbon atoms, which is optionally substituted by phenyl,

R³ denotes straight-chain or branched alkyl having up to 5 carbon atoms or a group of the formula —OR⁶,

wherein

R⁶ denotes hydrogen, a hydroxyl-protecting group or straight-chain or branched alkyl having up to 5 carbon atoms,

R⁴ denotes straight-chain or branched alkyl having 2 to 10 carbon atoms, which is optionally substituted by phenyl,

L denotes a radical of the formula —CO—, —CH(OH)—, —CH₂—, —CH(N₃)— or —CH(OSO₂R⁷)—,

wherein

R⁷ denotes straight-chain or branched alkyl having up to 4 carbon atoms or phenyl,

R⁵ denotes straight-chain or branched alkyl having 3 to 8 carbon atoms which is substituted by phenyl, or denotes benzyl or 2-phenylethyl,

D represents hydrogen, or represents a group of the formula —SO₂—NR⁸R⁹,







wherein

R⁸ and R⁹ are identical or different and denote hydrogen, phenyl or straight-chain or branched alkyl having up to 6 carbon atoms, which is optionally substituted by hydroxyl, or, together with the nitrogen atom, form a 5- to 6-membered saturated heterocyclic radical which has up to 2 further hetero atoms from the series consisting of S, N and/or O and is optionally substituted, including via a free N function, by straight-chain or branched alkyl having up to 6 carbon atoms, which in turn can be substituted by hydroxyl, and


E represents straight-chain or branched alkyl having up to 8 carbon atoms, and tautomers and salts thereof.

-78-

Preferred compounds include:



U.S. Patent No. 5,294,612 discloses compounds of the formula



wherein:

R^1 is hydrogen, alkyl, C_4 to C_7 cycloalkyl, C_4 to C_7 cycloalkyl substituted by C_1 to C_{10} alkyl or hydroxyl, 2- or 3-tetrahydrofuranyl, 3-tetrahydrothienyl 1,1, -dioxide, C_4 to C_7 cycloalkyl- C_1 to C_{10} alkyl, carboxy- C_1 to C_{10} alkyl, carbo- C_1 to C_4 lower-alkoxy- C_1 to C_{10} alkyl, dialkylamino C_1 to C_{10} alkyl, phenyl- C_1 to C_4 lower-alkyl, phenyl- C_1 to C_4 lower-alkyl in which the phenyl ring is substituted in the 2, 3, or 4-position by one or two substituents, the same or different, selected from the group consisting of amino, halogen, C_1 to C_{10} alkyl, carboxyl, carbo- C_1 to C_4 lower-alkoxy, carbamoyl, $NHSO_2$ -(quinolinyl), nitro and cyano;


R^3 is, C_1 to C_4 lower-alkyl, phenyl- C_1 to C_4 lower-alkyl, lower-alkoxyphenyl- C_1 to C_4 lower-alkyl, di- C_1 to C_4 lower-alkoxy-phenyl- C_1 to C_4 lower-alkyl, pyridyl- C_1 to C_4 lower-alkyl, C_4 to C_7 cycloalkyl- C_1 to C_4 lower-alkyl, phenylamino, di- C_1 to C_{10} alkylamino, halogen, trifluoromethyl, C_1 to C_4 lower-alkylthio, -cyano or nitro; and

R^6 is a nine or ten membered bicyclic ring having carbon and from one to two nitrogen atoms, and the heterocycle is made up of fused 5 or 6 membered rings or such ring substituted at any available carbon atom by one or two substituents, the same or different, selected from the group consisting of C_1 to C_4 lower-alkyl, halogen, C_1 to C_4 lower-alkoxy, C_4 to C_7 cycloalkyloxy, 4-morpholinyl, C_1 to C_4 lower-alkoxy- C_1 to C_4 lower-alkoxy, hydroxy, imidazolyl, oxo and 4-morpholinyl- C_1 to C_4 lower-alkoxy, or at any available nitrogen atom by C_1 to C_4 lower-alkyl, C_2 to C_4 lower-alkanoyl, or trifluoroacetyl; or a pharmaceutically acceptable acid-addition salt thereof.

Preferred compounds include:

1-Cyclopentyl-3-methyl-6-(4-quinolinyl)-
pyrazolo[3,4-d]pyrimidin-4-one

WO 93/12095 discloses compounds of the formula



or a pharmaceutically acceptable salt thereof,

wherein R^1 is H, C_1-C_4 alkyl, C_1-C_4 alkoxy or $CONR^5R^6$;

R^2 is H or C_1-C_4 alkyl;

R^3 is C_2-C_4 alkyl;

R^4 is H, C_2-C_4 alkanoyl optionally substituted with NR^7R^8 , (hydroxy) C_2-C_4 alkyl optionally substituted with NR^7R^8 , $CH=CHCO_2R^9$, $CH=CHCONR^7R^8$, $CH_2CH_2CO_2R^9$, $CH_2CH_2CONR^7R^8$, $SO_2NR^7R^8$, $SO_2NH(CH_2)_nNR^7R^8$ or imidazolyl;

R^5 and R^6 are each independently H or C_1-C_4 alkyl;

R^7 and R^8 are each independently H or C_1-C_4 alkyl, or together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, morpholino or 4-(NR^{10})-1-piperazinyl group wherein any of said groups is optionally substituted with $CONR^5R^6$;

R^9 is H or C_1-C_4 alkyl;

R^{10} is H, C_1-C_3 alkyl or (hydroxy) C_2-C_3 alkyl;


and n is 2, 3 or 4;

with the proviso that R^4 is not H when R^1 is H, C_1-C_4 alkyl or C_1-C_4 alkoxy.

Preferred compounds include:

2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]phenyl}-8-methylquinazolin-4-(3H)-one;
 2-{5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]-2-n-propoxyphenyl}-8-methylquinazolin-4-(3H)-one;
 8-methyl-2-{5-[2-(4-methyl-1-piperazinylcarbonyl)-ethenyl]-2-n-propoxyphenyl}quinazolin-4-(3H)-one;
 8-carbamoyl-2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]phenyl}quinazolin-4-(3H)-one;
 and 8-ethylcarbamoyl-2-(2-n-propoxyphenyl)quinazolin-4-(3H)-one;
 and pharmaceutically acceptable salts thereof.

WO 93/07149 discloses compounds of the formula



or a pharmaceutically acceptable salt thereof,

wherein R^1 is C_1-C_4 alkyl;

R^2 is H, methyl or ethyl;

R^3 is C_2-C_4 alkyl;

R^4 is C_1-C_4 alkyl optionally substituted with NR^5R^6 , CN, $CONR^5R^6$ or CO_2R^7 ; C_2-C_4 alkenyl optionally substituted with CN, $CONR^5R^6$ or CO_2R^7 ; C_2-C_4 alkanoyl optionally substituted with NR^5R^6 ; $SO_2NR^5R^6$; $CONR^5R^6$; CO_2R^7 ; or halo;
 R^5 and R^6 are each independently H or C_1-C_4 alkyl, or together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, morpholino, 4-(NR^8)-1-piperazinyl or 1-imidazolyl group wherein said group is optionally substituted by one or two C_1-C_4 alkyl groups;

R^7 is H or C_1-C_4 alkyl;

and R^8 is H, C_1-C_3 alkyl or hydroxy C_2-C_3 alkyl.

Preferred compounds include:

6-(5-bromo-2-n-propoxyphenyl)-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-(5-morpholinosulphonyl-2-n-propoxyphenyl)-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

6-[5-(2-carboxyvinyl)-2-n-propoxyphenyl]-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;


6-[5-(2-t-butoxycarbonylvinyl)-2-n-propoxyphenyl]-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-[5-(2-morpholinocarbonylvinyl)-2-n-propoxyphenyl]-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

and 3-methyl-6-[5-(2-morpholinocarbonylethyl)-2-n-propoxyphenyl]-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

and pharmaceutically acceptable salts thereof.

European published patent application No. 0607439 discloses compounds of the formula




[in formula (1), ring A represents a benzene ring, a pyridine ring or a cyclohexane ring; ring B represents a pyridine ring, a pyrimidine ring, or an imidazole ring.

Provided that the ring A and the ring B are combined sharing two atoms and the atoms shared may be either a carbon atom or a nitrogen atom.

In the case where the ring A is a pyridine ring and that except the case where the ring B shares the nitrogen atom of this pyridine ring to combine therewith, the ring A is represented by


-83-



R¹, R², R³ and R⁴, each of which may be the same or different from one another, represent each a hydrogen atom, a halogen atom, a lower alkyl group which may be substituted with a halogen atom, a cycloalkyl group which may be substituted, a lower alkoxy group, a hydroxyalkyl group, a nitro group, a cyano group, an acylamino group, a carboxyl group which may be protected, a group represented by the formula



(wherein R⁷ represents a lower alkyl group, and n represents 0 or an integer of 1 to 2), or a group represented by the formula




(wherein R⁴⁵ and R⁴⁶, each of which may be the same or different from each other, represent each a hydrogen atom or a lower alkyl group; or R⁴⁵ and R⁴⁶ can form a ring which may contain another nitrogen atom or oxygen atom together with the nitrogen atom to which they are bonded with the proviso that this ring may be substituted); or, two of R¹, R², R³ and R⁴ may together form methylenedioxy, ethylenedioxy or a phenyl ring.

R⁵ represents a hydrogen atom, a halogen atom, a hydroxyl group, a hydrazino group, a lower alkyl group, a cycloalkyl group which may be substituted, a lower alkoxy group, a lower alkenyl group, a carboxyalkyl group which may be protected, a carboxyalkenyl group which may be protected, a hydroxyalkyl group, a carboxyl group which may be protected, a group represented by the formula



(wherein R⁸ represents a lower alkyl group, and m represents 0 or an integer of 1 to 2), a group represented by the formula -O-R⁹ (wherein R⁹ represents a hydroxyalkyl group which may be protected, a carboxyalkyl group which may be protected or a benzyl group which may be substituted), a group represented by the formula




(wherein R²³ represents a hydroxyl group, a lower alkyl group, a lower alkoxy group, a hydroxyalkyl group or a hydroxyalkoxy group), a heteroaryl group which may be substituted, a 1,3-benzodioxolyl group which may be substituted, a 1,4-benzodioxyl group which may be substituted, a 1,3-benzodioxolylalkyl group which may be substituted, a 1,4-benzodioxylalkyl group which may be substituted, a group represented by the formula -C(R²⁴)=X [wherein X represents an oxygen atom, a sulfur atom or a group represented by the formula =N-R¹⁰ (wherein R¹⁰ represents a hydroxyl group, a cyano group or a carboxyalkoxy group which may be protected); and R²⁴ represents a hydrogen atom or a lower alkyl group], or a group represented by the formula -NR¹¹R¹² (wherein R¹¹ and R¹², each of which may


-84-

be the same or different from each other, represent each a hydrogen atom, a lower alkyl group, a hydroxyalkyl group, an aminoalkyl group, a carboxyalkyl group which may be protected, an alkylcarbamoyl group, a carboxyalkylcarbamoyl group which may be protected, a heteroarylalkyl group which may be substituted, a 1,3-benzoxolylalkyl group or a 1,4-benzodioxylalkyl group; or, further, R^{11} and R^{12} can form a ring which may contain another nitrogen atom or oxygen atom together with a nitrogen atom to which they are bonded with the proviso that this ring may be substituted).


R^5 represents a hydrogen atom, a halogen atom, a hydroxyl group, an amino group, a lower alkyl group, a lower alkoxy group, a lower alkenyl group, a 1,3-benzodioxolylalkoxy group, a 1,4-benzodioxylalkoxy group, a phenylalkoxy group which may be substituted, a group represented by the formula




(wherein R^{13} and R^{14} , each of which may be the same or different from each other, represent each a hydrogen atom, a lower alkyl group or a lower alkoxy group; or, further, R^{13} and R^{14} may together form methylenedioxy or ethylenedioxy), a group represented by the formula




a group represented by the formula



a group represented by the formula




a group represented by the formula




(in these formulas, R^{15} and R^{16} , each of which may be the same or different from each other, represent each a hydrogen atom, a lower alkyl group or a lower alkoxy group; or, further, R^{15} and R^{16} may together form methylenedioxy or ethylenedioxy), a piperidine-4-spiro-2'-dioxan-1-yl group, a group represented by the formula


-85-



(wherein R^{48} and R^{49} , each of which may be the same or different from each other, represent each a hydrogen atom, a lower alkyl group or a lower alkoxy group; or, further, R^{48} and R^{49} may together form methylenedioxy or ethylenedioxy; and Z represents a sulfur atom or an oxygen atom), a group represented by the formula



(wherein R^{50} represents a hydroxyl group, a halogen atom, a lower alkyl group, a lower alkoxy group, a carboxyl group which may be protected, a cyano group, a hydroxyalkyl group or a carboxyalkyl group), a group represented by the formula




(wherein R^{17} represents a hydrogen atom, a lower alkyl group, an acyl group, a lower alkoxyalkyl group, a carboxyalkyl group which may be protected or a hydroxyalkyl group; Y represents a group represented by the formula $-(CH_2)_q-$ (wherein q is 0 or an integer of 1 to 8), or a group represented by

the formula




further, in the group represented by the formula $-(CH_2)_q-$, when q is an integer of 1 to 8, each carbon atom may have 1 to 2 substituent(s); and R^{18} represents a hydrogen atom, a hydroxyl group, a carboxyl group which may be protected, a cyano group, an acyl group, a heteroaryl group which may be substituted or a cycloalkyl group which may be substituted], or a group represented by the formula



(wherein R^{19} represents a hydrogen atom, a lower alkyl group, a lower alkoxyalkyl group, an acyl group, a carboxyalkyl group which may be protected or a hydroxyalkyl group; R^{20} , R^{21} and R^{22} , each of which may be the same or different from one another, represent each a hydrogen atom, a halogen atom, a hydroxyl group, an amino group, a nitro group, a lower alkyl group, a lower alkoxy group, a lower alkoxyalkyl group, a lower alkenyl group, an acyl group, an acylamino group, an alkylsulfonylamino group, a hydroxyiminoalkyl group, an alkylloxycarbonylamino group, an alkylloxycarbonyloxy group or a heteroaryl group which may be substituted; or, further, two of R^{20} , R^{21} and R^{22} may together form a saturated or unsaturated ring which may contain a nitrogen atom, a sulfur atom or an oxygen atom; and r represents 0 or an integer of 1 to 8)].

-86-

WO 93/06104 discloses compounds of the formula



or a pharmaceutically acceptable salt thereof,
 wherein R^1 is methyl or ethyl;
 R^2 is ethyl or n-propyl;
 and R^3 and R^4 are each independently H, or C_1-C_6
 alkyl optionally substituted with C_1-C_7
 cycloalkyl or with morpholino.

Preferred compounds include:

5-[2-ethoxy-5-(3-morpholinopropylsulphamoyl)-
 phenyl]-1,3-dimethyl-1,6-dihydro-7H-pyrazolo[4,3-d]-
 pyrimidin-7-one;


1-ethyl-5-[5-(n-hexylsulphamoyl)-2-n-propoxy-
 phenyl]-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-
 d]pyrimidin-7-one;

1-ethyl-5-(5-diethylsulphamoyl-2-n-propoxy-
 phenyl)-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-d]-
 pyrimidin-7-one;

and 5-[5-(N-cyclohexylmethyl-N-methylsulphamoyl)-2-n-
 propoxyphenyl]-1-ethyl-3-methyl-1,6-dihydro-7H-
 pyrazolo[4,3-d]pyrimidin-7-one;

and pharmaceutically acceptable salts thereof.

U.S. Patent No. 5,346,901 discloses compounds of the formula



wherein

R¹ is H, C₁-C₃ alkyl, C₃-C₅ cycloalkyl or C₁-C₃ perfluoroalkyl;

R² is H, C₁-C₆ alkyl optionally substituted by OH, C₁-C₃ alkoxy or C₃-C₆ cycloalkyl, or C₁-C₃ perfluoroalkyl;


R³ is C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₆ alkynyl, C₃-C₇ cycloalkyl, C₁-C₆ perfluoroalkyl or (C₁-C₆ cycloalkyl)C₁-C₆ alkyl;

R⁴ taken together with the nitrogen atom to which it is attached completes a pyrrolidinyl, piperidino, or morpholino group;

R⁵ is H, C₁-C₄ alkyl, C₁-C₃ alkoxy, NR⁷R⁸, or CONR⁷R⁸;

R⁷ and R⁸ are each independently H, C₁-C₄ alkyl, (C₁-C₃ alkoxy)C₂-C₄ alkyl or hydroxy C₂-C₄ alkyl; and pharmaceutically acceptable salts thereof.

European published patent application No. 0442204 discloses compounds of the formula



or a pharmaceutically acceptable salt thereof, wherein

R¹ is C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₅ cycloalkyl, C₁-₆ alkyl, or C₁-₆ alkyl substituted by 1 to 6 fluoro groups;

R² is C₁-₆ alkylthio, C₁-₆ alkylsulphonyl, C₁-₆ alkoxy, hydroxy, hydrogen, hydrazino, C₁-₆ alkyl, phenyl, -NHCOR³ wherein R³ is hydrogen or C₁-₆ alkyl, or -NR⁴R⁵, wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, hexahydroazepino, morpholino or piperazino ring, or R⁴ and R⁵ are independently hydrogen, C₂-₆ cycloalkyl or C₁-₆ alkyl which is optionally substituted by -CF₃, phenyl, -S(O)ₙC₁-₆ alkyl wherein


n is 0, 1 or 2, -OR⁶, -CO₂R⁷ or -NR⁸R⁹ wherein R⁶ to R⁹ are independently hydrogen or C₁-₆ alkyl, pro-

vided that the carbon atom adjacent to the nitrogen atom is not substituted by said -S(O)ₙC₁-₆ alkyl, -OR⁶ or -NR⁸R⁹ groups;


R is halo, C₁-₄ alkyl, C₁-₄ alkoxy, cyano, -CONR¹⁰R¹¹, CO₂R¹², C₁-₄ alkylS(O)ₙ, -NO₂, -NH₂, -NHCOR¹³ or SO₂NR¹⁴R¹⁵ wherein n is 0, 1 or 2 and R¹⁰ to R¹⁵ are independently hydrogen or C₁-₄ alkyl; and

-88-

(A) is a ring of sub-formula (a) or (b) :



(a)




(b)

Preferred compounds include:

2-(5-cyano-2-propoxyphenyl)-7-methylthiopyrimido-[4,5-d]pyrimidin-4(3H)-one,
 2-(5-carboxamido-2-propoxyphenyl)-7-methylthiopyrimido[4,5-d]pyrimidin-4(3H)-one, or
 2-(5-carboxamido-2-propoxyphenyl)-7-cyclopropylamino[4,5-d]pyrimidin-4(3H)-one,
 or a pharmaceutically acceptable salt thereof.

U.S. Patent No. 5,010,086 discloses compounds of the formula



wherein


R_1 and R_3 are hydrogen or lower-alkyl;
 R_2 is lower-alkyl or fluorinated lower-alkyl; and the
 pyridine-N-oxide is attached at the 4- or 3-position;
 or a pharmaceutically acceptable acid-addition salt
 thereof.

Preferred compounds include:

1,3-Dihydro-6-(4-pyridinyl)-5-trifluoromethyl-2H-
 imidazo[4,5-b]pyridin-2-one N-(py)-oxide

-89-

U.S. Patent No. 5,290,933 discloses compounds of the formula




or a pharmaceutically acceptable salt thereof, wherein R^1 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-5} cycloalkyl, C_{1-6} alkyl, phenyl, C_{1-6} alkyl or C_{1-6} alkyl substituted by 1 to 6 fluoro groups; and R^2 is hydrogen, $-NHCOR^3$, or $-CONR^4R^5$, wherein R^3 is C_{1-6} alkyl, R^4 is C_{1-6} alkyl and R^5 is hydrogen or C_{1-6} alkyl.

Preferred compounds include:

N-methyl 1,6-dihydro-6-oxo-2-(2-propoxyphenyl)-pyrimidine-5-carboxamide,
N,N-dimethyl 1,6-dihydro-6-oxo-2-(2-propoxyphenyl)-pyrimidine-5-carboxamide,
5-acetamido-2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
or
2-(2-propoxyphenyl)pyrimidin-4(3H)-one,
or a pharmaceutically acceptable salt thereof.

U.S. Patent No. 5,073,559 discloses compounds of the formula




or pharmaceutically acceptable salt thereof, wherein R^1 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-5} cycloalkyl, C_{1-6} alkyl, phenyl, C_{1-6} alkyl or C_{1-6} alkyl substituted by 1 to 6 fluoro groups; R^2 is hydrogen, hydroxy, C_{1-6} alkyl, phenyl, mercapto, C_{1-6} alkylthio, CF_3 or amino; R^3 is hydrogen, nitro, amino, C_{1-6} alkanoylamino, C_{1-6} alkoxy, C_{1-6} alkyl, halo, $SO_2NR^4R^5$, $CONR^4R^5$, cyano or C_{1-6} alkyl $S(O)_n$; R^4 and R^5 are independently hydrogen or C_{1-6} alkyl; and n is 0, 1 or 2; provided that R^3 is not hydrogen when R^1 is C_{1-6} alkyl or C_{2-6} alkenyl and R^2 is hydrogen or hydroxy.

Preferred compounds include:

2-(2-{2,2,2-trifluoroethoxy}phenyl)purin-6-one,
 2-(2-cyclopropylmethoxyphenyl)purin-6-one,
 2-(2-benzyloxyphenyl)purin-6-one,
 2-(2-propoxyphenyl)-8-trifluoromethylpurin-6-one,
 2-(2-propoxyphenyl)-8-phenylpurin-6-one,
 2-(2-propoxyphenyl)-8-methylpurin-6-one,
 2-(2-propoxyphenyl)-8-mercaptapurin-6-one,
 2-(2-propoxyphenyl)-8-methylthiopurin-6-one,
 2-(2-propoxyphenyl)-8-aminopurin-6-one,
 2-(2-propoxy-5-nitrophenyl)purin-6-one,
 2-(2-propoxy-5-aminophenyl)purin-6-one,
 2-(2-propoxy-5-acetamidophenyl)purin-6-one,
 2-(2-propoxy-4-methoxyphenyl)purin-6-one,
 2-(2-propoxy-5-methoxyphenyl)purin-6-one,
 2-(2-propoxy-4-methylphenyl)purin-6-one,
 2-(2-propoxy-5-fluorophenyl)purin-6-one,
 2-(2-propoxy-5-dimethylsulphamoylphenyl)purin-6-one,
 2-(2-propoxy-5-methylsulphamoylphenyl)purin-6-one,
 2-(2-propoxy-5-sulphamoylphenyl)purin-6-one,
 2-(2-propoxy-4-methylthiophenyl)purin-6-one,
 2-(2-propoxy-5-cyanophenyl)purin-6-one, and
 2-(2-propoxy-5-carbamoylphenyl)purin-6-one,
 or a pharmaceutically acceptable salt thereof.

International Patent Publication PCT/EP96/03024 (WO97/03675) discloses compounds of the formula:



and salts and solvates (e.g. hydrates) thereof, in which:

R^0 represents hydrogen, halogen or C_{1-6} alkyl;

R^1 represents hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl C_{1-3} alkyl, aryl C_{1-3} alkyl or heteroaryl C_{1-3} alkyl;

R^2 represents an optionally substituted monocyclic aromatic ring selected from benzene, thiophene, furan and pyridine or an optionally substituted bicyclic



ring attached to the rest of the molecule via one of the benzene ring carbon atoms and wherein the fused ring A is a 5- or 6-membered ring which may be saturated or partially or fully unsaturated and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulphur and nitrogen; and

R^3 represents hydrogen or C_{1-3} alkyl, or R^1 and R^3 together represent a 3- or 4- membered alkyl or alkenyl chain.

Preferred compounds include:

Cis-2,3,6,7,12,12a-hexahydro-2-butyl-6-(4-methylphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-isopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopentyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-cyclopropylmethyl-6-(4-methoxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3-chloro-4-methoxyphenyl)-2-methylpyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (5aR,12R,14aS)-1,2,3,5,6,11,12,14a-Octahydro-12-(3,4-methylenedioxyphenyl)-pyrrolo[1'',2'':4',5']pyrazino[2',1':6,1]pyrido[3,4-b]indole-5-1,4-dione;
 Cis-2,3,6,7,12,12a-hexahydro-2-cyclopropyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 (3S,6R,12aR)-2,3,6,7,12,12a-hexahydro-3-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione;
 and physiologically acceptable salts and solvates (e.g. hydrates) thereof.

The specific compounds of the invention are:

(6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole -1,4-dione (Compound A); and
 (3S,6R,12aR)-2,3,6,7,12,12a-hexahydro-2,3-dimethyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (Compound B);

and physiologically acceptable salts and solvates (e.g. hydrates) thereof.

Examples of cGMP PDE inhibitors contemplated in this invention are also described in United States Patent No. 5,346,901 and published International Patent Publication WO 94/28902, both of which documents are incorporated herein by reference.

Sildenafil, 1-[[3-(4,7-dihydro-1-methyl-7-oxo-3-propyl-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-4-ethoxyphenyl]sulfonyl]-4-methylpiperazine, and salts thereof are disclosed in WO 94/28902.

Phentolamine, 3-[[[(4,5-dihydro-1H-imidazol-2-yl)methyl](4-methylphenyl)amino]phenol, and salts and esters thereof, and the use of phentolamine in the treatment of sexual dysfunction is disclosed in United States Patent No. 5,731,339, also incorporated herein by reference.

Sildenafil and phentolamine are each known to treat sexual dysfunction. The effectiveness of phentolamine for treatment of sexual dysfunction is demonstrated by test procedures described in U.S. 5,731,339. Similar procedures can be used to determine the effectiveness of sildenafil and combinations of phentolamine and sildenafil.

Since the present invention relates to a method of treatment comprising the administration of a combination of two components, the components can be co-administered simultaneously or sequentially. Alternatively, a single pharmaceutical composition comprising sildenafil, or a pharmaceutically acceptable salt thereof, and phentolamine, or a

pharmaceutically acceptable salt or ester thereof, in a pharmaceutically acceptable carrier can be administered. The components of the combination can be administered individually or together in any conventional oral dosage form such as a capsule, tablet, chewable tablets, powder, cachet, suspension or solution. The formulations can be prepared using conventional pharmaceutical excipients and additives using conventional techniques. Such pharmaceutically acceptable excipients and additives include non-toxic compatible fillers, binders, disintegrants, buffers, preservatives, anti-oxidants, lubricants, flavorings, thickeners, coloring agents, emulsifiers and the like.

Information on formulations comprising sildenafil are disclosed in WO 94/28902. Representative formulations comprising phentolamine are disclosed in U.S. 5,731,339. It is contemplated that where the two active ingredients are administered as a single composition, the dosage forms as disclosed in the aforementioned patent or application may readily be modified using the knowledge of one skilled in the art.

A typical formulation for sildenafil comprises 25, 50 or 100 mg of active and as inactive ingredients, microcrystalline cellulose, anhydrous dibasic calcium phosphate, croscarmellose sodium, magnesium stearate, hydroxypropylmethylcellulose, titanium dioxide, lactose, triacetin, and FD&C Blue #2 aluminum lake.

A typical formulation for phentolamine is as follows:

Component	mg/Tablet (w/w%)
phentolamine mesylate, USP	40 (10)
Microcrystalline Cellulose, NF	341.6 (85.4)
Croscarmellose Sodium, NF	16 (4.0)
Colloidal Silicon Dioxide, NF	0.4 (0.1)
Magnesium Stearate, NF	2 (0.5)
Total	400 (100)

The following are exemplary formulations for the phentolamine mesylate/sildenafil citrate combination:

-94-

Direct Compression Formulation

Component	mg/Tablet
Phentolamine Mesylate	80
Sildenafil Citrate	100
Microcrystalline Cellulose	207.5-209.0
Croscarmellose Sodium	10
Silicon Dioxide	0.5
Magnesium Stearate	0.5-2
Total	400

The direct -compression formulation is manufactured by blending the active ingredients and excipients and compressing the mixture into tablets.

Wet-Granulation Formulation

Component	mg/Tablet
Phentolamine Mesylate	80
Sildenafil Citrate	100
Microcrystalline Cellulose	80
Lactose	114-115.5
Sodium Starch Glycolate	12
Povidone	12
Water	(evaporates)
Magnesium Stearate	0.5-2
Total	400

The wet-granulation formulation is manufactured using the following steps:

1. the active ingredients are combined with microcrystalline cellulose, lactose and sodium starch glycolate in a mixer/granulator;
2. povidone is added to water to form a solution;
3. the granulating solution (from step 2) is added to the powder blend (from step 1) with agitation to form a granulation, and the resulting granulation is dried;
4. the dry granulation is blended with magnesium stearate; and

-95-

5. the mixture is compressed into tablets.

Fast-Dissolving FormulationsA

Component	mg/Tablet
Phentolamine Mesylate	40
Sildenafil Citrate	50
Gelatin	30
Mannitol	29
Flavor	1
Water	(evaporates)
Total Dry Tablet Weight	150

The above tablet form is manufactured by:

1. forming a uniform dispersion achieved by adding the active ingredients and excipients to water with agitation;
2. filling aliquots of the dispersion into molds; and
3. lyophilizing to form dry tablets.

B

Component	mg/Tablet
Phentolamine Mesylate	40
Sildenafil Citrate	50
Microcrystalline Cellulose	95
Crospovidone	10
Sodium Bicarbonate	2
Citric Acid	2
Flavor	1
Total	200

The tablets are made by blending the combination of the actives and excipients and compressing the mixture into tablets.

The compounds in the combination of this invention for treating sexual dysfunction are administered in accordance with the treatment regimens described in each of the above listed publications. For example, for a combination of a Type V cGMP PDE inhibitors such as

-96-

Sildenafil in combination with phentolamine, the typical dosage is 5 to 100 mg of Sildenafil and 5 to 75 mg of phentolamine per dose, usually administered approximately one hour prior to intercourse. It is expected that the dosage of the individual components in the combination will be less than the dosage required when the individual components are administered alone. The exact dose of either component of the combination to be administered and the timing thereof is determined by the attending clinician and is dependent on the potency of the compound administered, the age, weight, condition and response of the patient. Where the components of a combination are administered separately, the separate dosage forms need not be administered simultaneously.

Since the present invention relates to treatment with a combination of active ingredients wherein said active ingredients may be administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form. That is, a kit is contemplated wherein two separate units are combined: for example, a sildenafil pharmaceutical composition and a phentolamine pharmaceutical composition. The kit will preferably include directions for the administration of the separate components. The kit form is particularly advantageous when the separate components must be administered in different dosage forms (e.g. tablet and capsule) or are administered at different dosage intervals.

What is claimed is:

1. A pharmaceutical composition for the treatment of human sexual dysfunction comprising a therapeutically effective amount of phentolamine or a pharmaceutically acceptable salt or solvate or ester thereof, a therapeutically effective amount of a cGMP PDE V inhibitor or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.
2. A composition of claim 1 wherein the cGMP PDE V inhibitor is sildenafil or a pharmaceutically acceptable salt or solvate thereof.
3. The composition of claim 1 wherein the phentolamine is phentolamine mesylate.
4. The composition of claim 1 wherein the sildenafil is sildenafil citrate.
5. The composition of claim 1 wherein the phentolamine is phentolamine mesylate and the cGMP PDE V inhibitor is sildenafil citrate.
6. A method of treating human sexual dysfunction comprising the simultaneous or sequential administration of a therapeutically effective amount of phentolamine or a pharmaceutically acceptable salt, solvate or ester thereof, and a therapeutically effective amount of a cGMP PDE V inhibitor or a pharmaceutically acceptable salt thereof.
7. The method of claim 6 wherein the cGMP PDE V inhibitor is sildenafil or a pharmaceutically acceptable salt or solvate thereof.
8. The method of claim 6 wherein the phentolamine is phentolamine mesylate.
9. The method of claim 6 wherein the cGMP PDE V inhibitor is sildenafil citrate.

-98-

10. The method of claim 6 wherein the phentolamine is phentolamine mesylate and the cGMP PDE inhibitor V is sildenafil citrate.

11. A kit comprising in separate containers in a single package, pharmaceutical compositions for use in combination to treat sexual dysfunction which comprises in one container a therapeutically effective amount phentolamine or a pharmaceutically acceptable salt, solvate or ester thereof in a pharmaceutically acceptable carrier and in a second container a therapeutically effective amount of a cGMP PDE V inhibitor or a pharmaceutically acceptable salt of solvate thereof in a pharmaceutically acceptable carrier.

12. A pharmaceutical composition for the treatment of human sexual dysfunction comprising a therapeutically effective amount of a first vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof, a therapeutically effective amount of a second vasodilating agent or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.

13. The pharmaceutical composition of claim 12 wherein said first vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof is an adrenergic blocker.

14. The pharmaceutical composition of claim 13 wherein said adrenergic blocker is an alpha-adrenergic blocker.

15. The pharmaceutical composition of claim 14 wherein alpha adrenergic blocker is selected from the group consisting of an alpha1-adrenergic blocker, an alpha2-adrenergic blocker or both an alpha1-adrenergic blocker and an alpha2-adrenergic blocker.

16. The pharmaceutical composition of claim 12 wherein said second vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof is a cGMP PDE inhibitor.

17. The pharmaceutical composition of claim 12 wherein said first vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof is an adrenergic blocker and said second vasodilating agent

or a pharmaceutically acceptable salt or solvate or ester thereof is a cGMP PDE inhibitor.

18. The pharmaceutical composition of claim 17 wherein the adrenergic blocker is selected from the group consisting of phentolamine, phentolamine mesylate, phentolamine hydrochloride, phenoxybenazmine, tolazoline, dibenamine, yohimbine, terazosin, doxazosin and prazosin.

19. The pharmaceutical composition of claim 17 wherein the cGMP PDE inhibitor is a cGMP PDE V inhibitor.

20. The pharmaceutical composition of claim 17 wherein the cGMP PDE V inhibitor is selected from the group consisting of: sildenafil,

(6R, 12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (Compound A), and

(3S,6R,12aR)-2,3,6,7,12,12a-hexahydro-2,3-dimethyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (Compound B) or a pharmaceutically acceptable salt or solvate thereof.

21. A method of treating human sexual dysfunction comprising the simultaneous or sequential administration of a therapeutically effective amount of a therapeutically effective amount of a first vasodilating agent or a pharmaceutically acceptable salt or solvate or ester thereof, a therapeutically effective amount of a second vasodilating agent or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

International Application No
PC1/US 99/07046

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/415 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GOMAA A ET AL: "Topical treatment of erectile dysfunction: randomised double blind placebo controlled trial of cream containing aminophylline, isosorbide dinitrate, and co-dergocrine mesylate 'see comments!.' BMJ (CLINICAL RESEARCH ED.), (1996 JUN 15) 312 (7045) 1512-5. , XP002115285 abstract the whole document ---	12-15,21
P,X	SOLI M ET AL: "Vasoactive cocktails for erectile dysfunction: chemical stability of PGE1, papaverine and phentolamine." JOURNAL OF UROLOGY, (1998 AUG) 160 (2) 551-5. , XP002115286 abstract the whole document ---	12-15,21
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

14 September 1999

Date of mailing of the international search report

28/09/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Economou, D

INTERNATIONAL SEARCH REPORT

International Application No

PC1/US 99/07046

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CHAO R ET AL: "Experience with intracavernosal tri-mixture for the management of neurogenic erectile dysfunction." ARCHIVES OF PHYSICAL MEDICINE AND REHABILITATION, (1994 MAR) 75 (3) 276-8 , XP002115287 abstract page 277, left-hand column, paragraph 4 - right-hand column, paragraph 3</p>	12-15,21
X	<p>MIRONE V ET AL: "Ketanserin plus prostaglandin E1 (PGE-1) as intracavernosal therapy for patients with erectile dysfunction unresponsive to PGE-1 alone." BRITISH JOURNAL OF UROLOGY, (1996 MAY) 77 (5) 736-9. , XP002115288 abstract page 737, right-hand column, paragraph 4 - page 738, left-hand column, paragraph 3 page 736, left-hand column, line 1 - right-hand column, paragraph 2</p>	12-15,21
X	<p>BENNETT A H ET AL: "An improved vasoactive drug combination for a pharmacological erection program." JOURNAL OF UROLOGY, (1991 DEC) 146 (6) 1564-5. , XP002115289 the whole document</p>	12-15,21
X,Y	<p>US 5 731 339 A (ZONAGEN, INC.) 24 March 1998 (1998-03-24) cited in the application column 3, line 45 - column 17, line 18 claims 1-37</p>	1-21
X,Y	<p>WO 94 28902 A (PFIZER, LTD.) 22 December 1994 (1994-12-22) cited in the application the whole document</p>	1-21
X,Y	<p>WO 97 03675 A (LABORATOIRE GLAXO WELLCOME S.A.) 6 February 1997 (1997-02-06) cited in the application the whole document</p>	1-21
X	<p>EP 0 611 248 A (B.M.R.A. CO. B.V.) 17 August 1994 (1994-08-17) the whole document</p>	12-15,21
Y		16-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/US 99/07046

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5731339 A	24-03-1998	AU 5576896 A	18-11-1996
		BG 102010 A	30-04-1998
		CA 2219502 A	31-10-1996
		CZ 9703393 A	18-03-1998
		EP 0767660 A	16-04-1997
		HU 9802825 A	28-06-1999
		LT 97168 A,B	25-06-1998
		LV 12038 A	20-05-1998
		LV 12038 B	20-08-1998
		MD 980007 A	31-07-1999
		NO 974965 A	23-12-1997
		NZ 307020 A	29-06-1999
		PL 323087 A	02-03-1998
		SI 9620058 A	30-06-1998
		SK 145897 A	03-06-1998
WO 9428902 A	22-12-1994	WO 9633705 A	31-10-1996
		ZA 9603380 A	08-11-1996
		AT 163852 T	15-03-1998
		AU 676571 B	13-03-1997
		AU 6797394 A	03-01-1995
		CA 2163446 A,C	22-12-1994
		CN 1124926 A	19-06-1996
		CZ 9503242 A	17-07-1996
		DE 69408981 D	16-04-1998
		DE 69408981 T	02-07-1998
		DK 702555 T	06-04-1998
		EP 0702555 A	27-03-1996
		ES 2113656 T	01-05-1998
		FI 955911 A	08-12-1995
		GR 3026520 T	31-07-1998
WO 9703675 A	06-02-1997	IL 109873 A	27-12-1998
		IL 121836 A	27-12-1998
		JP 9503996 T	22-04-1997
		LV 12269 A	20-05-1999
		NO 954757 A	24-11-1995
		NZ 266463 A	24-03-1997
		PL 311948 A	18-03-1996
		ZA 9404018 A	08-12-1995
		AU 704955 B	13-05-1999
		AU 6419196 A	18-02-1997
		BR 9609758 A	26-01-1999
		CA 2226784 A	06-02-1997
		CN 1195290 A	07-10-1998
		CZ 9800033 A	13-05-1998
		EP 0839040 A	06-05-1998
EP 0611248 A	17-08-1994	HU 9900065 A	28-05-1999
		NO 980153 A	10-03-1998
		PL 324495 A	25-05-1998
		SK 3998 A	08-07-1998
		US 5567706 A	22-10-1996